









Authors

Dr Katie Featherstone,

Reader, School of Healthcare Sciences, Cardiff University

Dr Andy Northcott,

Lecturer, School of Allied Health Sciences, De Montford University et al.









Authors:

Dr Katie Featherstone, Reader, School of Healthcare Sciences, Cardiff University featherstonek@cardiff.ac.uk (PI)

Dr Andy Northcott, Lecturer, School of Allied Health Sciences, De Montfort University. andy.northcott@dmu.ac.uk

Co-authors and contributors

Professor Jackie Bridges, School of Healthcare Sciences, Southampton University (Jackie.Bridges@soton.ac.uk)

Karen Harrison Denning, Consultant Admiral Nurse and Director of Research and Evaluation, Dementia UK (karen.harrison-dening@dementiauk.org)

Dr Jane Harden, School of Healthcare Sciences, Cardiff University (hardenj@cardiff.ac.uk)

Professor Sue Bale, Aneurin Bevan University Health Board (Sue.Bale@wales.nhs.uk)

Professor Jane Hopkinson, School of Healthcare Sciences, Cardiff University (hopkinsonjb@cardiff.ac.uk)

Dr Rosie Tope, Carers Wales (rosie@herc.org.uk)

Dr Alexandra Hillman, School of Social Sciences, Cardiff University (hillmanae1@cardiff.ac.uk)

Ms Mandy King, Cardiff and Vale NHS Trust (Kinga1@cardiff.ac.uk)

Funding

The National Institute for Health Research Health Services and Delivery Research programme. Project: 13/10/80

Table of Contents

Authors:	i
Co-authors and contributors	i
Funding	i
Summary	iv
Scientific Summary	iv
Background	iv
Objectives	v
Methods	v
Results	v
Conclusions	vii
Introduction	8
Acute hospital care	10
Dementia and hospitalization	11
Impact on patient outcomes:	12
Examining culture and organization	13
Do we know anything about best practice in acute settings?	14
Conceptualising resistance and refusal of care	14
Locating Resistance and Refusal of Care in the Acute Setting	
Implications for our research strategy	
Research objectives	
Methodology	16
Data collection and analysis	
Data sources	
Ethnographic fieldwork	
Sampling	
Ethical approvals	
Modes of analysis and interpretation	
Study Results	23
Resistance: a routine part of everyday care	_
Physical Resistance: Getting out of bed, standing, walking	
Verbal Resistance: Shouting, anger and agitation	
Resistance towards the timetabled rounds of the ward	
Resistance to medicines and the medication round	
Personal care	37
Observation round	38
Removal of equipment: Treatments and Tubes	38
Removal of sheets and clothing	39
Language & performance: communicating the 'rules' of the ward in response to resistance	41
Use of multiple interactional approaches	41
Orienting to the reality of their situation	43
Clear instructions to be obeyed	44
Appeal to the necessity of the institution	44

NIHR HS&DR 13/10/80 Dr Katie Featherstone, Cardiff University

Negotiation and bargaining	45
Family and carers responses to resistance	46
The moral classifications of resistance	47
Judgements of individuals: resistance as identity	47
Resistance as deliberate and wilful	47
The emotional and somatic impacts of the organisation and timetables of the ward	48
Difficulties communicating care needs	49
Disorientation- place and time	51
Ward strategies of care in response to resistance	51
Cultures of containment and restraint	52
Outsourcing care and ward cultures of containment	55
Visibility and invisibility	57
Discussion	57
Conclusion: interventions and change	60
Summary	60
Consultation, raising awareness and informing wider user communities and publics about our	
findings	61
Collaboration with the arts	62
Dissemination to ward staff	63
Need for training	63
From 'bundles of care' to 'hacks'	63
On-line training in collaboration with Dementia UK	64
Carers perspectives	65
NHS Trusts	65
Developing no-cost interventions at ward level	65
Feasibility study: an experienced-based co-design framework	66
Acknowledgements	68
Defevences	6-

Summary

Up to half of all acute UK hospital beds are occupied by a person living with dementia at any one time, typically for a condition unrelated to their dementia. However, people living with dementia are highly vulnerable within the hospital setting: their health can significantly and suddenly worsen during an admission. Change is needed to improve the care for people living with dementia during an acute hospital admission.

This study aimed to find ways in which the delivery of nursing care could be adapted to improve the quality of care people living with dementia receive. To do this, we focussed on a common but poorly understood feature of caring for people living with dementia, the refusal and resistance of care. Over 18 months, we observed care within ten wards over five hospitals (two wards at each hospital) across England and Wales.

We found that people living with dementia resisted or refused care frequently while admitted to a hospital ward. Every person living with dementia that we observed resisted care to some degree at some point of their hospital stay. Nursing staff typically responded to this resistance as if it were a symptom of dementia. Our observations suggested this response could be problematic. Often resistance was in direct response to the care the person was receiving. By responding to resistance as a symptom of dementia, nursing staff could inadvertently exacerbate the resistance. Common responses included repeating instructions, raised voices, containment to beds and continued attempts to complete a task the patient had rejected. These responses could trigger further resistance, outright refusal of care and escalating cycles of resistance that caused stress and anxiety for the patient, their families and carers, and staff. Our findings identified a number of small changes to care which, with further study, may improve the acute care experience.

Scientific Summary

The goal of this in-depth ethnographic study was to identify ways in which the social organisation of nursing care and the interactional care processes at the bedside can be structured to improve the quality and humanity of care for people living with dementia who have been admitted to an acute hospital with a co-morbid condition.

Background

The acute hospital setting has become a key site of care for people living with dementia. The Department of Health recognises that as many as one in four acute hospital beds in the UK will be occupied by a person living with dementia at any given time. In some areas, these figures may be under-estimates, with some hospitals reporting that up to 50% of acute admissions may also have a diagnosis of dementia, while a significant number of other patients may be living with dementia but yet to receive formal diagnosis. Dementia as a condition is often thought of as something cared for first in the community, then later in specialist settings and long-term care, but the prominence of the acute hospital setting and its impact on people living with dementia cannot be ignored.

People living with dementia are a highly vulnerable group within the hospital setting and following an acute admission their functional abilities can deteriorate quickly and significantly. People living with dementia are more likely to experience a delayed discharge, are more likely to be re-admitted, and are at much higher risk of dying during an admission and in the months immediately after discharge, compared to people without a diagnosis of dementia or other cognitive impairment.

Although there is recognition that many hospitals have initiatives to improve the environment and support people living with dementia, there is an acknowledgement that hospitals are struggling to respond to their needs. However, prior to developing interventions and implementing policy solutions, detailed research is required to understand the role and needs of healthcare staff caring

for this patient population and to explore what constitutes 'good care' for people living with dementia within the acute setting.

Objectives

The focus of this study was to examine a common but poorly understood phenomena within the acute setting: refusal and resistance to care. Refusal of care (also labelled in the literature as resistance, resistive behaviours or rejection of care) is characterised as non-compliant behaviour in response to healthcare staff. People living with dementia may refuse medications, food and personal care as well as diagnostic or therapeutic procedures. Whilst there is already a large body of literature examining refusal of care within primary and long-term service provision, little is known about the social and organisational context of refusal of care and how clinical teams in acute hospitals respond when they believe a person living with dementia is refusing care.

In response, this in-depth ethnographic study examined the everyday work of nurses and healthcare assistants caring for people living with dementia within acute hospital wards. Our research questions were: How do ward staff respond to refusal and resistance to food, drink and medicines by people living with dementia being cared for on acute hospital wards, and what is the experiences of refusal of care from the perspective of patients and their carers?

Methods

Our approach to ethnography was informed by the symbolic interactionist research tradition, which aims to provide an interpretive understanding of the social world, with an emphasis on interaction, focusing on understanding how action and meaning are constructed within a setting. Our ethnographic approach enabled an in-depth evidence based analysis of everyday care and enabled us to understand how ward staff responded to the care needs of people living with dementia and to follow the consequences of their actions over time. Importantly, we also examined how ward staff accounted for and made sense of their responses to the care needs of people living with dementia in these contexts. Ethnography allowed us to examine these elements and, importantly, the interplay between them. This understanding was examined in the context of a narrative synthesis of the existing literature about refusal of care more widely.

This ethnography was carried out in 10 wards within 5 hospitals across England and Wales purposefully selected to represent a range of hospitals types, geographies and socio-economic catchments. Across these sites, 155 days of observational ethnographic fieldwork were carried out in areas of acute hospitals known to admit large numbers of people with dementia for acute conditions: Trauma & Orthopaedic wards and Medical Assessment Units (MAU or variants thereof). Approximately 600,000 words of observational fieldnotes were collected, written up, transcribed, cleaned and anonymised by the ethnographers (KF and AN). To provide a detailed contextual analysis of the events observed, the expertise involved, and the wider conditions of patient care, we also carried out ethnographic (during observation) interviews with ward staff and family carers (n=436) and detailed case studies (total: n=10) of people living with dementia, observing care at the bedside throughout their admission, and carrying out interviews with the patient, their carers, family members, and the ward staff caring for them.

Ethics Committee approval for the study was granted by the NHS Research Ethics Service via the Wales Research Ethics Committee 3 on 24th June 2015 (15/WA/0191). Substantial amendments to the study protocol were approved at a meeting of the Wales REC 3 committee on 10th December 2015.

Results

In the course of our analysis we treat the concept of resistance as emergent and 'in the making', which surfaced in the ways ward staff interpreted and responded to behaviours during the mundane

encounters, interactions and the everyday routine work of the ward. The actual act (of refusal, resistance or rejection) is always context bound and involves many factors (which we have tried to capture in our analysis below) and should not be seen as an isolated feature of a dementia diagnosis. Thus, rather than classifying these behaviours as distinct entities, we use the term 'resistance' to represent the range of responses people living with dementia have to the ways in which their care is delivered at the bedside and their wider experiences of their admission to an acute hospital ward. In turn, these responses were viewed by staff as problematic, difficult, signified a lack of capacity, and because they do not 'fit' the organisation and timetabled routines of the ward they must be overcome or managed. Specifically, our analysis identified:

- High levels of resistance, refusal and rejection of care amongst people living with dementia within acute hospital wards. In total, we identified 1,052 (T&O=523, MAU=529) incidents or episodes of resistance to care. We identified that every person living with dementia observed within an acute hospital ward resisted care at some point during their admission.
- Resistance could manifest in a number of ways (in order): (1) a person attempting to get out (and getting out) of bed, standing and walking around (this includes: wanting to go home, trying to leave the bay, ward or hospital; going to other rooms or areas of the ward/unit; pulling at the bed rail; approaching the nursing station); (2) verbal and physical cues (this includes: shouting, angry, agitated; crying, screaming, sounding distressed; asking to go home, see family or be taken somewhere; biting, pushing, pulling, spitting at; or holding on to staff, refusing to leave the ward); (3) not submitting to the timetabled rounds of the ward (this includes: mealtimes, food and drink, medication, personal care, observation rounds and examinations); (4) and the removal of equipment (this includes: pulling or removing IV and/or gastric tubes, dressings, catheters, nebulisers, cannulas, oxygen mask, and other medical equipment crucial to care; pulling sheets off the bed or removing clothing).
- The subsequent patterns of responses and interventions typically involved nurses and HCAs using multiple interactional approaches that combined highly repetitive language (orienting the person to reality, providing instructions to be obeyed, emphasising the necessity and requirements of the institution, negotiation and bargaining) with the performative (a focus on carrying out work on the body, completing essential care, and containment of the person at the bedside). These approaches created damaging cycles of stress for patients, families, and for ward staff.
- Ward staff typically interpreted resistance as a feature of a dementia diagnosis, which
 overshadowed the person, with longer-term, chronic and disruptive resistance viewed as a
 feature of their identity, which could become their principal identity in the context of the ward.
 However, resistance to care was typically a response to ward organisation and delivery of care
 and typically rational to that person's present ontology and perceptions.
- Resistance appeared to be a response and reaction to the impacts of admission on individuals.
 These were both emotional and somatic and included the difficulties communicating need, high levels of anxiety, and the unfamiliar environment of the ward, which could lead to disorientation, and in turn, this was associated with the fixed routines, timetables and the organisation and delivery of care. A key impact of this was the person becoming viewed within the ward as having increased dependency, which in turn, resulted in them loosing skills and independence.
- The completion of ward routines and timetables dominated shifts, which meant that staff did not consider it possible to focus on what appeared to be low level resistance that was not an immediate priority or risk, particularly in the context of pressing demands of the ward routines of personal care, bed making, observation and medication rounds. Often subtle signs could be identified in a patient's body language and changes in their behaviour that indicated resistance or the potential for resistance. However, it was unusual for ward teams to recognise these early signs, or feel able to prioritise responses to them.
- At the bedside, staff response to resistance to care was one of containment and restraint.

 Raising the side rails of the bed, or tucking bed sheets in tightly around the patient were both

common means to contain a patient within the bed. For those patients sitting at the bedside, the close placement of the mobile tray table, unreachable walking frames and technologies such as chair alarms were used to contain people and keep them sitting in their bedside chair. Clinical technologies, including continence technologies (particularly full continence pads, and using bed pans, and commodes at the bedside), medication and sedation, and tightly secured medical equipment also limited or restrained movement from the bedside. Importantly, these approaches to patient care and their containment at the bedside was both a response to resistance, but were also frequently the trigger of resistance or patient anxiety.

Conclusions

This in-depth ethnographic study has examined the everyday work of nurses and healthcare assistants caring for people living with dementia within acute hospital wards. We identified key responses to resistance to care at the organisational and interactional levels of care delivery at the bedside that had significant consequences for people living with dementia, their family carers, and also for ward staff. Our analysis, drawn from observations that allowed the ethnographers to spend extended time within and across shifts observing specific ward bays and the people within them. This observational approach revealed the potential underlying reasons or triggers for a person's resistance to care. In response to resistance, we identified powerful cultures of containment and restraint, with the interactional care work in the ward driven by the organisational demands of delivering care within fixed routines and timetables that do not meet the needs of people living with dementia. A key impact was the invisibility of people living with dementia and their care needs within acute wards.

At the bedside, staff management and response to resistance to care by people living with dementia was one of containment and restraint. Although specific techniques had some variance between wards, the overall strategy was always to keep the person living with dementia within their bed or sitting at the bedside. Across all sites, staff expressed high levels of concern and anxiety about people attempting to or leaving the bed or bedside, and this increased exponentially if they were walking in the bay, the wider ward and corridor or close to the ward entrance. Importantly, these approaches to patient care and their containment at the bedside was both a response to resistance, but were also frequently the trigger of resistance or cause of patient anxiety.

We identified that at an organisational level, a key response to resistance by people living with dementia within wards was to assign one-to-one agency HCA staff to care for them. This was a policy within almost all hospitals and wards. The outsourcing of dementia care and expertise via 'dementia workers' and one-to-one agency care meant that the care of people living with dementia was typically seen as other people's work and this could have powerful impacts on the wider ward culture. Ward staff typically interpreted resistance as a feature of a dementia diagnosis, which overshadowed the person, and could become their principal identity in the context of the ward. A key impact of this was the person becoming viewed within the ward as having increased dependency, which in turn, resulted in individuals loosing further skills and independence.

Importantly, our analysis has identified ways in which the social organisation of nursing care and the interactional care processes at the bedside can be structured to improve patient and family care experience and the effectiveness of treatments. In response, we are currently using our findings to develop simple no-cost innovations ('hacks') at the interactional and organisational level within wards that can be introduced and used by nurses and HCAs. We are also developing on-line awareness raising and training films tailored for staff within the acute setting in collaboration with Aneurin Bevan Health Board, Admiral Nurses, and Dementia UK. We are currently developing and testing the feasibility of these interventions within one acute 'laboratory' ward.

An Evidence based Investigation into the care that people living with dementia receive following an acute hospital admission

Introduction

The quality of care delivered to vulnerable older people admitted to acute hospital settings in England and Wales has been under the spotlight following the publication of both the Francis report (2013) and the Andrews report (2014). These enquiries identified unacceptable quality of care, including the systemic deprivation of dignity and respect (Francis report, 2013) and serious concerns about the culture of care (Andrews report, 2014). Such findings were not new, with a 2006 Joint Committee report concluding that 'an entire culture change is needed' (HL 156-I, HC 378-I 2006:3) to improve the care of people living with dementia receive in hospital "We believe that this question of the culture in a hospital is absolutely crucial" (HL 156-I, HC 378-I:44, 2006). Both the Joint Committee and the Andrews report gave particular focus on the poor support with medication, dehydration and malnutrition older people experience in hospital.

The Care Quality Commission has repeatedly identified systemic failures in the care of older people (2011) and concluded that the variation in care in hospitals experienced by people with dementia meant that they are 'likely to experience poor care at some point along their care pathway' (Care Quality Commission, 2014:9). A number of national audits have also been critical of the systematic failings in the quality of care people living with dementia receive in hospital (NAO, 2007) and of particular concern was the failure to safeguard people living with dementia from avoidable and protracted hospital stays (Health Foundation, 2011, section 5.9). Findings supported by the more recent national Adult Inpatient Survey, which concluded that older people experienced inconsistent or poor standards of dignity and respect, emphasizing that this was 'a significant general problem affecting inpatients in the vast majority of NHS acute hospital trusts' (Vizard and Burchardt, 2015:3). With variable or poor practice in the care of people living with dementia identified within 56% of hospitals inspected (Care Quality Commission, 2014).

There are corresponding widespread concerns amongst carers and families about the quality of hospital care people living with dementia experience (Patients Association, 2009), with the Carers Trust reporting hospitalisation leading to dehydration, sores and a lack of nutrition (Newbronner, et al 2013). The Nottingham City and Nottinghamshire County Joint Health Scrutiny Committee (2010) review identified carers concerns about mealtime support, medications, poor management of behaviours arising from anxiety, and low levels of staff awareness and understanding of how to care for people living with dementia.

In response to this body of evidence, there has long been recognition by policymakers of the potential for the improvement of care for people living with dementia in hospitals, particularly when they are admitted to general hospital for an unrelated condition (Health Foundation, 2011). A 'transformation of dementia services' has been called for within the Department of Health national strategy, 'Living Well with dementia' (Department of Health, 2009:14) and by the Dementia Action Alliance (2012). In partnership with the NHS Institute for Innovation and Improvement, a key objective within their Call to Action (2012) is for services to be designed around the person with dementia through the creation of dementia friendly hospitals. This focuses on five key areas: the care environment, knowledge, skills and attitudes of the workforce, the ability to identify and assess cognitive impairment, supportive discharge back home, and person-centred care plans involving families and carers. The Alzheimer's Society (2013) recommends hospitals endorse and implement these objectives.

NICE (2006) have previously recommended that acute and general hospital trusts should plan and provide services that address the specific personal and social care needs and the mental and physical health of people living with dementia who use acute hospital facilities. Improved screening, management and discharge processes are key to improving outcomes in the acute hospital (National Audit Office, 2007). The Department of Health have highlighted the need for individuals with direct responsibility for dementia within the acute setting and that these individuals should work closely with specialist older people's mental health teams (Department of Health 2013). More recently, the Prime Minister's Challenge on Dementia 2020 (Department of Health, 2015) renewed the focus on creating dementia-friendly health and care, with the goal of every person with dementia obtaining the safest, best care in our acute hospitals.

A large number of reports highlight that in order to deliver these significant improvements in care, there is a need for specialist training for staff to improve their knowledge, skills and attitudes within the acute setting (Royal College of Psychiatrists, 2006; NICE, 2006; Department of Health, 2009, National Audit Office, 2007; Department of Health and Social Care, 2011; Department of Health, 2013). Importantly, the majority of nursing staff (89%) have identified working with people living with dementia as challenging (Alzheimer's Society 2008), with healthcare professionals within the acute setting lacking the necessary skills and knowledge to care for this patient population (Scottish Government, 2010; Department of Health, 2009; Tadd et al, 2011). Without the appropriate training and support, there is a recognition that healthcare staff will become resentful, demoralised and cut themselves off from patients, all things that can lead to the de-personalisation and dehumanization of people in their care (Cornwell, 2012).

With the failure to provide appropriate training for hospital staff in caring for people living with dementia identified as a key contribution to their poor outcomes and long inpatient stays (Alzheimer's Society, 2007). Training in the care and support of people living with dementia should be part of the core curriculum for nurses, highlighting that such training must also be part of their continuous professional development (Department of Health, 2009), with a further emphasis on training required for all staff working in health or social care (Department of Health and Social care, 2011; Alzheimer's Society. 2013). Training is recognized as the key to reducing stigma (Department of Health, 2009) and delivering dignity in care (Department of Health (2015). The House of Lords House of Commons Joint Committee on Human Rights advocates training for hospital staff that utilises a human rights framework to improve the culture of dignity and respect for older people in hospital (HL 156-I, HC 378-I, 2006).

However, there is still a disparity between these policy recommendations and their implementation within the acute setting. Many hospitals now have initiatives to improve the environment and support they provide for people living with dementia (Alzheimer's Society, 2015), and there are examples of innovative practice (Prime Minister's Challenge, 2012), however, even within institutions where high quality acute care for people living with dementia is identified, this may be limited to specific wards and fail to reach across an organisation (Alzheimer's Society (2015). Overall, there is an acknowledgement that hospitals are struggling to cope with the challenge of an ageing population in the context of increasing hospital admissions for this group (Royal College of Physicians, 2013). Hospital doctors report high levels of concern about the lack of continuity of care for older patients (Royal College of Physicians, 2013).

Yet despite calls for a culture change across the NHS (HL 156-I, HC 378-I, 2006; Department of Health, 2015) and key national objectives and strategies to reduce variability and improve care, the detrimental impact of a hospital admission on the long-term independence of people living with dementia has been consistently documented. There is still unacceptable variation in the quality of care for people living with dementia in hospital and at discharge (Alzheimer's Society, 2015). The Care Quality Commission's most recent report (2017) suggests that although the provision of health

and social care for people living with dementia in some areas of England has moved away from a 'tipping point'(CQC, 2017:3) services within other geographic areas have moved closer to that point of crisis.

Acute hospital care

There is increasing recognition that a key aspect of care for people living with dementia that needs immediate action is improving their experiences and outcomes following an admission to hospital for an unrelated condition (Health Foundation, 2011; Department of Health, 2009; House of Commons Public Accounts Committee, 2008; the National Institute for Health and Clinical Excellence (NICE) and the Social Care Institute for Excellence (SCIE); 2006). The NAO similarly advocates increased investment to improve care for people living with dementia within general hospitals to enable long-term savings and shorter hospital admissions (NAO, 2007). The Alzheimer's Society most recent report (2016) has found that poor care for people living with dementia is still widespread, that the quality of care varies widely between hospitals, with the admission for a person living with dementia up to five to seven times longer than other patients over the age of 65 in the worst-performing hospitals (Alzheimer's Society, 2016).

UK health policy emphasizes the importance of keeping people living with dementia out of hospital, minimizing the duration of any necessary hospital admission and supporting people living with dementia in the community (NICE SCIE, 2006). However, although health and well-being outcomes for people living with dementia are better when they are supported within the community, there is also acknowledgment that in reality many people living with dementia also need to be cared for in hospitals and other institutions.

Hospitals are not designed to care for this group (Tolson et al., 1999). Evidence suggests that the structural conditions and standardized care plans within the acute setting often do not fit the needs of people living with dementia and their families (Pinkert and Holle, 2012), with one ethnographic study identifying systemic and organizational factors within the hospital setting compromised clinical staff's ability to provide dignified care (Tadd et al, 2011). Hospital systems are designed to care for patients with one clinical problem, but is a setting 'which is chiefly subscribed by people who have many things wrong' and who are thus deemed as 'inappropriate' (Rockwood & Hubbard, 2004:429) patients. This leads to the treatment of the acute clinical problem for patient admission being prioritized, with the care of their additional dementia not recognized by clinical staff as a priority (Tolson et al., 1999). However, older patients commonly have more than one chronic condition in addition to their acute illness (Tadd et al 2011), and for people living with dementia, there is a need for specific care practices that brings together care of dementia and their acute clinical problem with the acute care setting and its practices (Hart et al., 2002).

Within the acute setting there is often the perception that the person living with dementia is at fault rather than the environment (Kitwood, 1993). Clinical staff often hold negative stereotypical attitudes towards older patients in their care (Lothian and Philip, 2001), with clinical features of dementia such as 'wandering' perceived by clinical staff as a deviant behavior (Dawson and Reid, 1987; Radar, et al, 1985). This lack of understanding of the needs of people living with dementia often results in this group being labelled "difficult" (Moyle et al, 2008).

The care of patients living with dementia within hospitals is not only as a welfare issue but also as a human rights issue. The House of Lords House of Commons Joint Committee on Human Rights highlights concerns about poor treatment, neglect, abuse, discrimination (HL 156-I, HC 378-I, 2006: paragraphs 9 - 65) and the lack of dignity especially for personal care needs (HL 156-I, HC 378-I, 2006). The report 'Counting the cost of care' (Alzheimer's Society 2008) identified the person with dementia being treated with a lack of dignity and respect as a key area of. The Alzheimer's Society survey found that over one third (36%) of carers said the person living with dementia was never

treated with respect and dignity (Lakey, 2009). A more recent Alzheimer's Society report (2016), reported that 60% of carers reported experiencing a lack of dignity or understanding.

The undignified care people living with dementia experience in hospital does not happen in a vacuum; it is rooted in the wider social discrimination older people experience within our society (Age UK, 2012.). The Joint Committee on Human Rights identified the powerful impacts of historic and embedded ageism that have contributed to the systematic failures to respect and protect the human rights of older people within the healthcare system (HL 156-I, HC 378-I, 2006). This seems particularly notable in the acute setting, where discrimination in the provision of services and quality of care has been acknowledged as 'an ageist policy' (RCP, 2006:5). The Department of Health implementation plan for the older people's national Service framework (Philp, 2006), singled out the deep-rooted negative attitudes and behaviours towards older people in acute care.

This stigmatizing of older people in hospitals is particularly acute for people living with dementia (Bamford et al, 2004). The poor recognition and systematic under-treatment of post-operative pain amongst people living with dementia (Morrison and Sui, 2000; Banicek, 2010) 'illustrates the discriminatory care provided for people with dementia' (Department of Health, 2009:62). In addition, people living with dementia from black, Asian and minority ethnic (BAME) communities are more likely to experience stigma and poor care (House of Commons All-Party Parliamentary Group on Dementia, 2013), while women with dementia are exposed to a 'triple jeopardy' (Graham et al, 2003; Bamford, 2011) of discrimination because of their age, sex and condition.

There is evidence to suggest that clinical staff routinely believe that people living with dementia do not belong within the acute setting, which is seen as an inappropriate place for older people more generally to receive care (Tadd et al, 2011). However, there was little exploration by staff of where a more appropriate place for older people would be. Tadd et al suggest that this may indicate an underlying ageism (Tadd et al, 2011). But it can also be seen as an example of 'specialism hegemony' – acute staff tend to dismiss conditions they see as belonging to the specialism of mental health.

The Kings Fund report (2012) highlights that 'the work of caring for older patients with complex needs is hard and testing – physically, psychologically and emotionally'(p4). This work carried out by nurses and healthcare assistants is often described as 'basic' rather than 'essential' care, implying that this is unskilled work and the report acknowledges the physical and emotional impact of this work 'In hospital, patients with dementia and delirium may disturb other patients, or may be challenging and difficult to keep calm and safe...changing a doubly incontinent patient and remaking the whole bed requires the effort of two people. Sometimes patients are unresponsive, and beds need to be changed again almost immediately'(p4).

This presents a significant NHS challenge (Royal College of Physicians, 2013) and new approaches are needed for improving what happens to older people, and more specifically people living with dementia when they are admitted to hospital. Consumer groups have lobbied to improve the experience of elderly patients (Alzheimer's Society, 2008; Patients Association, 2010), a need also recognized by several government enquiries (National Confidential Enquiry into Patient Outcome and Deaths, 2009; Care Quality Commission, 2013). However, the research agenda has lagged behind (Royal College of Physicians, 2013) and there is an evidence vacuum in understanding the experience of elderly patients with dementia and how their care can be improved within the acute setting.

Dementia and hospitalization

The acute hospital setting has become a key site of care for people living with dementia. The Department of Health (DoH) recognises as many as one in four acute hospital beds in the UK will be

occupied by a person living with dementia at any given time (Alzheimer's Society, 2009; DoH 2014 cited in Alzheimer's Society 2016), the equivalent of 3.2 million bed days per year. In some areas, these figures may be under-estimates, with some hospitals reporting that up to 50% of their acute admissions may also have dementia (Alzheimers Society 2016). A systematic review has identified that internationally, the prevalence estimates for people living with dementia within acute hospitals range from between 12.9–63.0% across studies (Mukadam and Sampson, 2011). Dementia as a condition is often thought of as something cared for first in the community, then later in specialist settings and long-term care, but the prominence of people living with dementia in the acute hospital setting and the impact admissions have upon these patients cannot be ignored.

A diagnosis of dementia is associated with an increased risk of hospitalization (Phelan et al, 2012). Russ et al (2012) estimate that approximately 6% of all people living with dementia in the UK are inpatients in acute hospitals at any one time, in comparison to only 0.6% of people over 65 without a diagnosis of dementia. A hip fracture (Holmes, 1999; Pinkert and Holle, 2012), urinary tract infection (Sampson et al, 2009), pneumonia (Sampson et al, 2009), and nutritional disorders (Pinkert and Holle, 2012) are amongst the common causes of admission amongst people living with dementia. A national review of case notes for people living with dementia (7987) found the majority were admitted within care of the elderly (40%), general medical (25%) and orthopaedic (11%) acute care wards (Royal College of Psychiatrists, 2013).

Although prevalence rates differ by hospital and are dependent on their specific population, current estimates are believed to be low due to underreporting or late diagnosis of this population (NAO, 2007). Estimates suggest that within the acute setting, approximately 50% of those affected by dementia do not yet have a formal diagnosis in their medical records (Goldberg et al, 2012; Sampson et al, 2009; Russ et al, 2012). There are a range of potential reasons for this potential under-diagnosis, (Koch et al, 2010) or delayed diagnosis of dementia, with much of this due to clinical teams not having the appropriate expertise (Koch et al, 2010). A significant number of people living with dementia may have their first assessment when they are admitted with an acute condition (Holmes, 1999). For example, amongst older people admitted to hospital following a hip fracture, of those living with dementia (40%), just over a quarter (27%) received their diagnosis during their admission (Holmes, 1999).

However, there are also likely to be many older people within the acute setting who may appear to who have features of dementia, but who may have different underlying causes of cognitive decline including delirium or sub-syndromal delirium. With one screening study of a large cohort of older patients following an unplanned admission within an acute hospital setting (MAU) found a high prevalence of delirium (15.5%), and a high rate of undiagnosed (72%) delirium amongst this population (Collins et al, 2010). With other studies identifying similar high levels of co-morbid mental health in this group (Cooper, 2010; Goldberg et al, 2012). Medication and co-morbid chronic conditions such as diabetes can also impact on cognitive function (Russ et al, 2012) within the acute setting.

Impact on patient outcomes:

Yet people living with dementia are a highly vulnerable group within this setting and following an acute admission the functional abilities of this population can deteriorate significantly (Goldberg et al, 2012). A systematic review has identified that people living with dementia in the acute hospital are older, require more hours of nursing care, have longer admissions, at higher risk of delayed discharge and likely to experience functional decline during their admission (Mukadam and Sampson, 2011). Similarly, a screening study of emergency admissions of patients over 70 years old with cognitive decline, the majority were at risk of malnutrition (80%) and just under half (47%) were classified as incontinent and needed help at meal times (49%) (Phelan et al, 2012). With an acute hospital admission for people living with dementia associated with an increased risk of

functional decline (Creditor, 1993; Sager et al, 1996) and of experiencing adverse events (Creditor, 1993; Watkin et al, 2012). With a longitudinal cohort study of acute emergency admissions found that patients over 70 with cognitive impairment had markedly higher short-term mortality, with 24% dying during admission (Sampson et al, 2009). With this group more likely to be re-admitted and also more likely to die post-admission than patients admitted for the same reason (Care Quality Commission, 2014). Similarly, a prospective cohort study, found that people living with dementia who has an unplanned acute hospital admission had half the survival time of similar inpatients without a diagnosis of dementia (Sampson, et al, 2013).

The poor recognition and systematic under-treatment of pain amongst people living with dementia (Morrison and Sui, 2000; Banicek, 2010) illustrates the inequity in care experienced by people living with dementia in the acute setting (Department of Health, 2009: p62). People living with dementia receive poor end-of-life care, fewer palliative medications (Sampson et al, 2006), and only a third of level of painkilling opioid medication (Morrison and Sui, 2000) compared to patients without a diagnosis of dementia. Pain is poorly identified and undertreated in people living with dementia (Sampson et al, 2015). People living with dementia find it difficult to articulate their pain (Banicek, 2010), with one study finding that they receive only a third of the opioid medication provided to other patients who do not have a diagnosis of dementia, concluding that the majority of people living with dementia were in severe pain post-operatively (Morrison and Sui, 2000; Sampson, et al, 2015).

People living with dementia in the acute setting are also at risk of 'cascade iatrogenesis', where the treatment or intervention used to treat the initial acute admitting condition, leads to an unintended sequence of multiple medical complications and a cascade of decline in the person (Thornlow et al, 2009) which can result in further dependency, institutionalisation, and potentially death during their acute admission (George et al, 2013). Adverse events are common for people living with dementia during an acute admission and importantly, these are typically associated with identifiable risk factors, with staff failing to understand and recognise the impact of a diagnosis of dementia on patient outcomes (Watkin et al, 2012).

George et al (2013) in their review of challenges and solutions to the care of people living with dementia in the acute setting conclude that "We need to increase awareness and understanding of the ways in which the manifestations of healthcare related harm are different in these patients, often presenting as the geriatric syndromes such as falls, delirium, incontinence and functional decline. The subtlety of presentation and detection of adverse events in patients with dementia is compounded by a culture of 'low expectation', which runs counter to the required 'safety culture' (George et al, 2013: 360).

Examining culture and organization

The House of Lords concluded that 'an entire culture change is needed' (HL 156-I, HC 378-I:3) to improve the care of people living with dementia receive in hospital "We believe that this question of the culture in a hospital is absolutely crucial" (HL 156-I, HC 378-I:44).

Research highlights the importance of the culture of care in influencing frontline delivery. Metaethnography findings (Bridges et al, 2013) reflect that, in spite of acute care nurses' aspirations to deliver a high standard of psychosocial care, this was largely dependent on ward-level social and organisational conditions. NIHR studies of acute settings report a strong association between ward climate and quality of care. Patterson et al. (2011) found that more positive patient and carer assessments of acute care for older patients was correlated with higher staff ratings of team climate in terms of "supporting each other", highlighting the vital role of the ward manager in shaping a positive climate for care. Findings mirrored in a second SDO study (Maben et al. 2012) that the experiences of working in wards impacts directly on patient experiences of care and meant that complex or demanding patients received less personalised care from staff (Maben et al, 2012).

Ward nurses playing a key role in facilitating the flexibility needed for hospital systems and processes to function effectively (Allen and Lyne, 1997) SDO studies have examined the role and impact of AHPs (Petchey et al, 2013) and Advanced Nursing Practitioners (Williamson et al, 2012) in ward leadership. Studies highlight the importance of focussing on the relationships between nurses and less privileged groups, which influences how care is organized, supervised and delivered (Daykin and Clarke, 2000) given the increasing delegation of "hands-on" care to HCAs (Daykin and Clarke, 2000).

Do we know anything about best practice in acute settings?

Few studies have explored what constitutes 'good care' for people living with dementia within the acute setting, with studies developing models of care for this patient group typically assuming that the principles of good care that have been integrated into long-term facilities, can transfer unproblematically into other settings (Moyle et al, 2008). Only a small number of studies have evaluated proposed models of care to improve patient outcomes within acute settings for dementia, however, the focus of these interventions has been to address 'chronic confusion', (c.f. Inouye et al., 1999; Li et al., 2003; Bradley et al., 2005) rather than dementia. There is little evidence that training interventions to enhance healthcare worker expertise and capacity in working specifically with patients with dementia are effective. Recent reviews (Elliott et al, 2012; Moyle et al, 2008) suggest that prior to developing interventions, further detailed research to understand the role and needs of healthcare workers with this patient population (Elliott, et al, 2012) and exploring what constitutes 'good care' within the acute setting (Moyle et al, 2008) is needed.

In their review identifying best practice for caring for this patient population within the acute setting, Moyle et al (2008) identified a number of key features: the early identification and assessment, the knowledge and attitudes of nursing staff, a multidisciplinary approach to care, providing focused communication, a reduction in stressors, carer and family involvement. Staff within an Australian hospital reported that the main constraints in being able to provide best practice were environmental, sociocultural, and economic, concluding that this setting was not appropriate for this patient group (Borbasi et al, 2006).

Conceptualising resistance and refusal of care

Resistance and refusal of care by people living with dementia is not a clearly defined concept within the wider research literature. The terms 'resistance' and 'refusal' are used interchangeably, both clinically and in the literature, to cover both the range and extent of this behaviour or phenomena (Ishii et al 2012). Other terms used in the literature include 'non-compliant behaviour' (Kable et al, 2012), 'challenging behaviours' (Cutson et al, 2016) 'behaviours that challenge' (Keady and Jones, 2010), and as 'agitated' (Logsdon et al, 1998) and 'aggressive' (Stevenson et al, 2009) behaviours. At their core, this remains something of an umbrella term and covers behaviour(s) characterized as non-compliance in response to healthcare (Kable et al, 2012).

Within the research examining what Ishii et all (2012) conceptualize in their review under the umbrella term 'rejection of care', there is a significant focus on long-term settings including community care, home care, care home and nursing home settings, psychiatric wards, and specialist mental health settings. Within these settings there has been sufficient research to support systematic reviews that examine how dementia and associated behaviours, including rejection of care, affect: friendship (Harris 2012), marriage (Baikie 2002), caregivers (Bonda et al, 2003; Torti et al, 2004; Schoenmakers, et al, 2010; Seitz et al, 2012) the person as patient (De Boer et al, 2007; Seitz et al, 2012), self and identity (Caddell and Linda, 2010), quality of life (Beerens, et al. 2013), the costs of care (Quentin, et al. 2010; Knapp et al, 2012; Jonsson and Anders 2009), access to care

(Cooper et al. 2010; Mukadam, et al 2011), care staff resilience (Elliott, et al, 2012), care staff stress (Pitfield et al. 2011), care staff competence (Traynor et al, 2011), care staff communication skills (Eggenberger et al, 2012; Vasse et al. 2010) and care staff retention (Chenoweth et al. 2010). The impact of behaviours related to a dementia diagnosis, including rejection of care, have been explored extensively within long-term and specialist settings.

Locating Resistance and Refusal of Care in the Acute Setting

Thus, while there is considerable evidence exploring the long-term settings in which one would expect to find people living with dementia being cared for, research examining the impact of other settings where people living with dementia also belong is scarce. A notable setting where there is little research is the acute care setting, where people living with dementia form a significant population. Acute care nurses in the UK have reported that they are 'always' responsible for caring for at least one patient with a diagnosis of dementia (Dewing and Dijk 2014) and almost all find such care a 'challenging' part of their work (Alzheimer's Society 2008). 'Rejection of care' is a recognised feature of these admissions (Werner at el 2002, Moyle et al 2008, 2010, Tadd et al 2011; Dewing and Dijk, 2014), yet unlike other long-term and specialist settings, few studies have examined the prevalence, causes, experiences and impacts of this phenomenon within the acute ward.

There remains a significant gap in the literature that provides systematic evidence of the extent of refusal and resistance to care, how it manifests, the ways in which ward staff respond, and whether cultures of restraint continue to exist in acute settings in response to rejection of care behaviours, and why. The limited literature examining rejection of care in the acute hospital setting acknowledges this, highlighting the scarcity of high quality studies examining this phenomena within the acute environment (Werner et al 2002, Barratt 2004, Rantala et al 2014 Pizzacalla et al 2015, Cheong et al 2016).

Implications for our research strategy

This small body of research highlights that there is a clear need for systematic research exploring the ways in which resistance and refusal manifests in the acute care setting, and what can be done to improve care in response to the phenomena to enhance both care experiences and outcomes.

To do this we must first address the tension that lies in the clear articulation of what we mean and understand by resistance and refusal when observing it. In other words, knowing it when we see it. Thus, a key aspect of our examination and analysis of resistance, refusal and rejection of care is by focusing on the ways in which ward staff (nurses and HCAs) interpret, recognize, assess, classify, and subsequently respond to these behaviours. Ishii et al (2012) give a very helpful definition of 'rejection of care': to "reject evaluation or care (e.g. bloodwork, taking medications, ADL assistance) that is necessary to achieve the patient's goals for health and well-being" (p12). Importantly, they go on to state that this definition requires a patient's *intent* to refuse or reject care and does not constitute a definition if the patient's somnolence renders them unable to swallow medication. This definition also excludes aggressive behaviours. Yet as we will show, within our study, a person being non-responsive or aggressive in response to care delivery were key features of how ward staff practically recognised and understood dementia, with 'refusal of care' interpreted within the ward as an expected feature or outcome of the behavioural or psychological symptoms of dementia during an admission.

In the course of our analysis we treat the concept of resistance as emergent and 'in the making', which surfaced in the ways staff interpreted and responded to behaviours during the mundane encounters, interactions and the everyday routine work of the ward. The actual act (of refusal, resistance or rejection) is always context bound and involves many factors (which we have tried to capture in our analysis below) and should not be seen as an isolated aspect within a wider repertoire of dementia-related behaviours. Thus, rather than classifying these behaviours as distinct entities,

we use the term 'resistance' and see the ways in which these manifest as part of a continuum of responses people living with dementia have to the ways in which their care is delivered at the bedside and their wider experiences of their admission to an acute hospital ward. In turn, these responses are viewed by staff as problematic, difficult, believed to signify a lack of capacity, and because they do not 'fit' the organisation and timetabled routines of the ward they must be overcome or managed.

Research objectives

In response, this study used an in-depth ethnographic approach to examine the work of nurses and healthcare assistants who are responsible for refusal of care with a focus on mealtimes and medications within wards to explore how staff respond when people living with dementia refuse care. We have focussed on wards known to have a large number of people with cognitive impairment (Medical Admissions Unit and Orthopaedic wards) within a purposive sample of 5 hospitals in England and Wales. Ethnography is particularly useful when exploring complex and sensitive topics in health care. Our analysis has focused on identify ways in which the social organisation of nursing care and care processes can be structured to support care to improve patient and family care experience and the effectiveness of treatments.

Methodology

Ethnography can provide sophisticated tools for understanding the complexities of the everyday and examining daily meaning-making (Ybema et al, 2009) within an organisational setting. Ethnography allows a detailed understanding of organisational culture, organisational change, and the interrelationships between different elements of an organisation. Importantly, it also takes into account the perspectives of patient, carer, clinical team and wider hospital staff (Hammersley, 2006; Caracelli, 2006). It is particularly useful to examine research questions and topics where measurement is either not easy or inappropriate, where the aim is to access the unspoken and tacitly understood, and where the topic is complex and highly sensitive (Dixon-Woods, 2003).

Our approach to ethnography is informed by the symbolic interactionist tradition, which aims to provide an interpretive understanding of the social world, with an emphasis on interaction. This approach focuses on understanding how action and meaning are constructed within a specific setting and acknowledges the mutual creation of knowledge by the researchers and the researched (Guba and Lincoln, 1994). Thus, within this study, the aim of our approach is to uncover the relevant conditions of people living with dementia within the acute hospital setting and to understand how the wide range of social actors within these settings (the large number of ward staff they will come into contact with during their admission) actively respond to these conditions through their actions, and the consequences of their actions. Ethnography allows us to examine these elements, but importantly, the interplay between them. It examines 'up close and in person' how work is organized and how the organizing organizes people' (Ybema, et al, 2009:1).

Ethnography is the in-depth study of a small number of cases. Ethnographers study people's actions and accounts within their natural everyday settings and collect relatively 'unstructured' data from a range of sources including observation, informal interviews, and documentary evidence (Hammersley and Atkinson, 1989). It is also important to consider the articulation work of people within organisational and institutional settings, examining how people within them account for and make sense of their actions. Ethnographers 'hold that an appreciation of the extraordinary-in-theordinary may help to understand the ambiguities and obscurities of social life' (Ybema et al, 2009:2). The value of this approach is the depth of understanding and theory generation it can provide, with a key objective to provide findings which are transferable to other settings (Hammersley, 1987).

The aim is to explore the details of everyday life that can otherwise go unnoticed, trying to read the tacitly known scripts and schemas that organize ordinary activities. Starr (1999) notes the importance of examining organizational infrastructure and the 'hidden mechanisms' (p377) within them that are constructed and embedded into the technical and procedural work carried out within it. It also examines, not just the front stage performance, but also the backstage work practices (Goffman, 1959). Within any organization there are also always groups whose everyday work is not recognised formally and is often unnoticed and invisible (Starr, 1999). In the hospital setting this includes carers, nurses, healthcare assistants, cleaners and porters. In the context of understanding how healthcare services within hospital settings are delivered and the organisation underlying its delivery, ethnography can examine how the social and institutional forces shape and influence the work of health care providers (Greenhalgh and Swinglehurst, 2011) and the everyday routine behaviours of individuals, both within and across multi- disciplinary teams (Quinlan, 2009).

There is a long tradition of using ethnography within healthcare settings (Higginbottom et al, 2013). There are many examples of studies that provide detailed ethnographic findings that have had significant impact on policy (Popay and Williams, 1998:34). Ethnography's potential to inform public debate, policy and practice is increasingly recognized (Lauder, Brown and Halsey 2004; Hammersley 2004). There has also been a growing legitimation of qualitative evidence as appropriate to health services research and the EBM movement (Popay and Williams, 1998; Black, 1994). In terms of the presentation of findings, ethnographic 'thick description' provides the reader with ways to connect concepts, policies and practice to detailed empirical examples (Vaughan 2005). These details allow the reader to develop not only a strong connection to that social world but also an understanding of the complex social relations in the context of both the personal impact and how it connects with wider public and policy issues (Vaughan 2005).

Thus, this study has focussed on the largely invisible routine work of caring for people living with dementia, each being significant elements of the everyday routine care carried out by nurses and HCAs in acute hospital wards. We provide a detailed understanding of the social and institutional forces that shape and influence this work. Our ethnographic approach has enabled us to understand how staff respond to the care needs of people living with dementia and to follow the consequences of their actions. Importantly, we have also examined how they account for and make sense of their responses to the care needs of people living with dementia in these contexts. Ethnography allows us to examine these elements and, importantly, the interplay between them (Atkinson et al, 2008).

Data collection and analysis

Data collection (observations and interviews) and analysis has been informed by the analytic tradition of grounded theory (Glaser and Strauss, 1967), a practical and flexible approach for ethnographic research (Charmaz and Mitchell, 2001). It uses the constant comparative method and theoretical sampling whereby data collection (observation and interview data) and analysis are interrelated (Glaser and Strauss, 1967; Corbin and Strauss, 1990) and carried out concurrently (Green, 1998; Suddaby, 2006). The flexible nature of this approach is important, because it has allowed us to increase the 'analytic incisiveness' (Charmaz and Mitchell, 2001:160) of the study: as data was collected in one site, preliminary analysis of this was proceed in parallel, with this preliminary analysis informing the focus of later stages of data collection and analysis.

Whilst these traditions have developed independently, they are complementary and grounded theory strengthens the ethnographic aims of achieving a theoretical interpretation of the data, whilst the ethnographic approach prevents grounded theory from being applied in a mechanistic and rigid way (Charmaz and Mitchell, 2001). A common concern with an ethnographic approach is that it can treat everything within a setting as data, which can lead to the ethnographer collecting large volumes of unconnected data and producing a heavily descriptive analysis (Coffey and Atkinson, 1996). This approach provides a middle ground in which the ethnographer, often seen as

a passive observer of the social world, can use grounded theory to provide a systematic approach to data collection that can be used to develop theory to address the interpretive realities of the range of actors within this setting (Charmaz and Mitchell, 2001).

Data sources

This ethnography was carried out in 10 wards within 5 hospitals across England and Wales purposefully selected to represent a range of hospitals types, geographies and socio-economic catchments. Across these sites, 155 days of observational ethnographic fieldwork were carried out in areas of acute hospitals known to admit large numbers of people with dementia for acute conditions: Trauma & Orthopaedic wards and Medical Assessment Units (MAU or variants thereof). Approximately 600,000 words of observational fieldnotes were collected, written up, transcribed, cleaned and anonymised by the ethnographers (KF and AN). To provide a detailed contextual analysis of the events observed, the expertise involved, and the wider conditions of patient care, we also carried out ethnographic (during observation) interviews with ward staff and family carers (n=436) and detailed case studies (total: n=10) of people living with dementia, observing care at the bedside throughout their admission, and carrying out interviews with the patient, their carers, family members, and the ward staff caring for them.

Ethnographic fieldwork

Within each hospital setting we conducted non-participant observation over a 4-6 week period carried out in areas of acute hospitals known to admit large numbers of people with dementia for acute conditions (MAU and Orthopaedic wards), with a focus on the everyday work of nurses and HCAs to examine the everyday clinical and care processes and responses to refusal and resistance to care. Approximately 600,000 words of observational fieldnotes that have been written up, transcribed, cleaned and anonymised (by KF and AN).

Multi-sited ethnography defines the object of study via a number of techniques or tracking strategies and within the fieldwork we recognise the importance of focussing on the 'busy intersections' (Rosaldo, 1989:28) and of seeking out sites of tension where a large number of interests and identities are expressed. It is argued that it is at these points that identity and culture become articulated, enacted and constructed. Our aim was to provide a detailed understanding of the clinical and interactional work and processes that influence nursing, HCAs and other clinical staff (this included mealtime assistants, SPRs, consultants, AHPs, and staff with managerial responsibilities) response to 'refusal of care' with a focus on everyday care, medication rounds and meal times. We have studied actions and accounts within their natural everyday settings to explore how individuals, teams, wards, and hospitals, respond to and manage refusal and what influences these approaches. Our focused observational strategy within each setting:

- Concentrated on the work of nurses and HCAs and other clinical staff from a range of
 disciplines and roles when they are involved in the care of people living with dementia,
 focusing on medication rounds and meal times. We have mapped the organisation of care,
 responses to refusal, management, and communication of refusal of care with wards.
- Followed nurses and HCAs within each ward setting to explore their everyday work, and
 what informs this work. Identifying the processes of decision-making, the management of
 uncertainty, and treatment procedures in response to refusal of care. Examining the
 everyday routine behaviours of individuals and within wards.
- Focussed on observing handovers, admissions, and conversations with carers, which are all
 opportunities for sharing information about refusal behaviours and how these might best
 be managed.

• Where possible, collected routine data (from ward managers and within patient records) about ward staffing levels, overall work allocation, bed occupancy, patient acuity, turnover and the recorded levels of resistance and refusal at the time of fieldwork to provide context and an understanding of the workload of resistance and refusal within this setting.

This has provided a detailed understanding of organisational and care processes that impact on the responses to and the management of refusal and resistance within this patient group. We examined the everyday work of staff, their practices and the interactions between staff and with patients and carers within these wards.

Ethnographic interviews with patients living with dementia and their carers:

Where possible, we carried out ethnographic (during observation) interviews with patients and their carers within the ward setting to explore the experiences and needs of this patient population and issues of refusal from their perspectives. While we originally proposed to interview patients resisting care, this was ultimately not possible as we underestimated the emotional duress and agitation linked to such experiences. Where possible we spoke to people living with dementia following a period of resistance, but the prior resistance was not raised to them, in order to minimise further agitation.

Thus, in response to these limitations, in consultation with ward staff and the NHS REC committee, we used our case studies to obtain the perspectives of people living with dementia and their families and carers via ethnographic interviews. Ethnographic interviews were carried out with the case study participants (n=71) and their family members (n=37), to explore:

- Experiences of admission and care, what is the impact of the physical environment and wider hospital structures
- Being listened to, communication and decision-making
- Fears and concerns about treatment and management, particularly around refusal and resistive behaviours
- Identification of refusal of care and involvement and recognition of carer expertise

Ethnographic interviews with nurses and HCAs:

To provide a detailed understanding of the influences on healthcare professionals response to 'refusal of care'. Ethnographic (during observation) interviews have been carried out with nursing, HCAs and clinical staff from a range of disciplines (this may include, mealtime assistants, SPRs, consultants, AHPs, and staff with managerial responsibilities) (n= 436) as they are caring for this patient group within each ward with a focus on medication rounds and meal times. This allowed us to question what they are doing and why:

- What is the articulation work within those settings, how do staff account for and make sense of their actions?
- What is the experience and training of working with people living with dementia and refusal of care, what informs their practice?
- What aspects of caring are defined as 'difficult', demanding or rewarding and what is their confidence in competence with working with this patient group. What are the barriers and enablers to supporting this patient group?
- What is the recognition and rewards of providing care for this patient group from patients, relatives, colleagues, managers?

Case studies:

Across the hospital sites we carried out detailed case studies (total: n=10) of patients living with dementia identified as refusing or resisting food or medicines (target: n=5). This has allowed us to

extend our fieldwork to provide a detailed contextual analysis of the events, the clinical staff and expertise involved and the wider conditions of their care and their impacts over time. This provides understandings of the broader care systems within the acute setting that impact on their care and provide a multi-perspective analyses. Purposive sampling was used, informed by our early analysis of observational data within each setting. Each case study has involved:

- Detailed systematic observations of patient care during their admission (114 days of observation), interviews with carers, and family members (n=37), and where possible, patients (total: 71), to explore the needs of this patient population and issues of refusal (Target= 20-40).
- We have also carried out interviews with the nurses and HCAs and other clinical staff (this
 included, feeding assistants, SPRs, consultants, AHPs, and staff with managerial
 responsibilities) involved in the care of these patients (see number of ward staff interviews
 above) to explore their response to refusal or resistive behavior with a focus on medication
 rounds and meal times (see interviews above).

This has allowed us to follow the impact of the everyday routine care carried out by nurses HCAs and other clinical staff and the consequences of their response and management of refusal and resistive behavior for this patient group and their carers.

Field notes of observation and near verbatim text have been written up into word files (Van Maanen, 2011; Emerson, et al., 2011) and all audio recordings of observations and interviews (ethnographic and in-depth) have been written up in word files or transcribed verbatim by a professional transcription service. All sites, individuals, and data collected has been anonymised and sorted in line with the Data Protection Act 1998, and NHS England Data Protection Policy 2014. Storage of the data is managed by the Cardiff University Information Security Framework Program.

Sampling

Sampling in ethnography requires a flexible, pragmatic approach, using a range of variables that may influence the phenomena, and what is known based on the available literature. Probability sampling is not appropriate, instead non-probability sampling, which is not representative of the wider population was used to provide analytically rather than statistically generalizable findings (Curtis et al, 2000; Mays and Pope, 2000). This is the most appropriate to study organisations or a clearly defined group and the size of the sample required for this approach is determined by the nature and scope of the study aims. Using this approach, the number of sites and participants in the sample is judged appropriate not on the basis of size, but on the quality and appropriateness of the sample and when saturation of data has been achieved (Mays and Pope, 2000).

Setting and access of hospitals:

This approach emphasizes the importance of comparisons across sites (Vogt, 2002) allowing for and optimizing the generalizability of findings (Herriott and Firestone, 1983) and enhancing the ability for the findings to impact on policy and practice (Caracelli, 2006). Hospital settings are well suited to an ethnographic approach. At first glance, hospitals may appear to operate in similar ways, however, they often have their own unique culture informed by local dominant cultures and belief systems, which in turn means that care and decision making can vary widely within institutions (Van Der Geest and Finkler, 2004; Goodson and Vassar, 2011). Thus, we identified a range of variables that may influence the phenomena using purposive and maximum variation sampling to include 5 hospitals that represent hospitals types, geographical location, expertise, interventions and quality (Marshall, 1996). Our 5 acute hospital settings have been identified from across the UK to represent the:

 Types of acute hospital (2 large University teaching hospitals (B and D), 2 medium sized general hospitals (A and C) and 1 smaller general hospital (E))

- Geographical locations to include urban (A), inner-city (C, D) and a large rural and urban catchment area (B, E), situated across England and Wales.
- A range of specialist and non-specialist clinical and non-clinical staff from no formal expertise to dementia specialist workers (B).

Sampling within each hospital site: Whilst our data collection sites (acute hospitals) are standardized, with sequentially and systematic data collection, there was some variation within each site. We have used theoretical sampling within sites to ensure that representativeness and consistency of concepts and events is achieved within the study, rather than sites and people. Informed by grounded theory, sensitizing concepts from the ongoing analysis informed the next stage of data collection to expand the research process and to capture relevant aspects as they emerged into the ongoing analysis. The focus was on 'discovery' to ensure the grounding of emerging concepts within data and the reality of the settings (Glaser and Strauss, 1967).

Sampling of wards for observation:

We observed episodes of care involving patients within the MAU and Orthopaedic wards which receive a large population of patients who have dementia, who require acute medical attention: MAU (Medical Assessment Unit): This is where unscheduled admissions arrive for assessment from A&E, the outpatient department clinic or their GP. Following assessment patients are discharged, transferred to a specialist centre, or admitted to an inpatient bed (Collins et al, 2010). These are high turnover settings, with the goal of discharging or transferring patients within 24 hours. There is no routine within this setting, with staff geared to deal with acute admissions with fast turnaround, with no continuity or personalised care, and a chaotic atmosphere. Thus, they are not an area designed or conducive to patients who have any cognitive deficit or dementia at a critical time where escalation of symptoms may occur. Our Carer Steering Group all had poor experiences of this setting and found it a frightening time, where they felt they were not listened to, and were often separated from their partner (n=529).

Orthopaedic wards: These settings allowed us to observe routine ward care, and patients who have an unscheduled admission following an accident where there may be no opportunity to provide additional support. The population of patients within this setting have had an accident or fall that has typically resulted in a fracture (n=65).

Sampling and recruitment of staff for observation and interviews:

We followed the work of nurses and HCAs. We purposively sampled to ensure that across the ward settings we included the range of clinical grades (clinical support worker nursing, nurse associate, entry level nurse, nurse specialist, nurse team leader, advanced nurse, nurse team manager, modern matron, nurse consultant) and other clinical staff (this included mealtime assistants, SPRs, consultants, AHPs, and staff with managerial responsibilities) responses to 'refusal of care'. Within each acute setting we worked with our key contact, who was typically the senior nurse responsible for care of the elderly and people living with dementia within each trust, who facilitated the process of identifying and introducing the team to key informants at hospital and ward levels. These key informants also provided introductions to our wards.

Sampling and recruitment of patients and carers for interview and observation:

Within wards we focussed on capturing the "incidents, events, and happenings that denote the work that they do, the conditions that facilitate, interrupt, or prevent their work, the action/interaction by which it is expressed and the consequences that result" (Corbin and Strauss, 1990: 421). Importantly, the focus of observation was on the daily practice of nurses and HCAs and

other clinical staff during mealtimes and medication. It was not possible to predict the patients and carers within each hospital ward during the fieldwork period, however we found that a high number of patients in the wards we observed had a diagnosis of dementia (as identified in ward records).

Case studies:

We carried out a focused series of individual case studies (n=10), identified using purposive sampling informed by our early analysis of our initial observational data. We include a range of subjects (maximum variation sampling) who have had particular types of experiences within the setting (critical case sampling). Thus, sampling included patients who represented a range of presenting, diagnostic and prognostic factors, and where possible, socio-demographic factors:

- We expected to identify refusal and resistance via the ward team and ward records, however, although we found high rates of refusal in the wards, we found low rates of identification, recognition and recording.
- Men (n=5) and women (n=5), aged over 65 (70-99 years old) with unplanned admission (fractured hip = 5, fractured wrist= 1, high number of falls = 2; head injury= 1; gout = 1) and an accompanying co-morbid diagnosis of dementia (including: Dementia = 6, '?Dementia'= 1, Alzheimer's= 1, and vascular dementia= 2) formally recorded in their medical records.
- Expected length of stay. This can be highly variable for this group, from days to weeks and months, thus we followed these patients for up to 6 weeks within each hospital setting.
- Where possible we carried out follow-up interviews with individuals and families after their discharge. Of these patients, two died during their admission, six were medically fit and waiting for a placement in a nursing home, and two were medically fit and waiting for a placement in a care home.

Ethical approvals

Ethics Committee approval for the study was granted by the NHS Research Ethics Service via the Wales Research Ethics Committee 3 on 24th June 2015 (15/WA/0191). Substantial amendments to the study protocol were approved at a meeting of the Wales REC 3 committee on 10th December 2015. The committee has approved this research project for the purposes of the Mental Capacity Act 2005 and confirms that it meets the requirements of section 31 of the Act in relation to research carried out as part of this project on, or in relation to, a person who lacks capacity to consent to taking part in the project. The study was accepted by NHS Research Permissions Wales on 16th July 2015, with NIHR CSP and West Midlands CRN on 11th March 2016 and with the Health Research Authority on 27th May 2016. Recruitment for the study was managed and recorded through the Central Portfolio Management System and closed on 31st January 2017.

Modes of analysis and interpretation

Data collection (observations and interviews) and analysis has been informed by the analytic tradition of grounded theory (Glaser and Strauss, 1967). We have utilized the constant comparative method and theoretical sampling whereby data collection (observation and interview data) and analysis are interrelated (Glaser and Strauss, 1967; Corbin and Strauss, 1990) and are carried out concurrently (Green, 1998; Suddaby, 2006). The flexible nature of this approach is important, because it allowed us to increase the 'analytic incisiveness' (Charmaz and Mitchell, 2001:160) of the ethnography. As these data were collected in one site at a time, preliminary analysis proceeded in parallel with data collection, with this preliminary analysis informing the focus of later phases of data collection in the next site and the further subsequent stages of analysis.

We applied an inductive approach to our analysis, a widely used approach, which means we developed our hypothesis from the data, rather than a priori (Pope et al, 2000). Analysis involved

the development and testing of analytic concepts and categories, and the strategies we used for their development, included the careful reading of the data, looking for patterns and relationships, noting anything surprising and inconsistencies and contradictions across the range of perspectives gathered. Initially this produced a collection of 'sensitizing concepts' (Blumer, 1954) and analytic memos, which then informed the development of more refined and stable analytic concepts. Line-by-line coding is not appropriate for fieldnotes, instead, coding was selective and involved whole events or scenarios (Charmaz and Mitchell, 2001). The constant comparative method means that the coding of data into categories was a recurrent process. The data was then examined in the context of previous fieldwork and the analytic memos generated then informed further data collection within the next site and the next, more focused, stages of analysis (Charmaz and Mitchell, 2001). The analytic concepts that emerged from this process were then further tested and refined to develop stable concepts that we believe transcend the local contexts to identify broader structural conditions (Corbin and Strauss, 1990) influencing responses to refusal of care.

A key aspect of our approach to analysis was to also take a systematic approach to the development of the ethnographic analysis into tangible outputs and interventions, via the following phases: (1) We have explicitly utilised multiple perspectives (sociological, policy, clinical, patient and carer) to inform the development of our analysis to identify both local organisational insights and broader structural conditions (Corbin and strauss, 1990), which we believed might influence care. The analysis of the observational data was supplemented by, and triangulated with, data from clinical, carer, and patient interviews. (2) The development of our analysis into ward based outputs was always in close collaboration with our Carer Steering Group and our participating hospital sites and ward teams to establish relevance and feasibility within wards, and their utility and transferability to other acute hospital settings that have different organisational contexts (Pawson &Tilley, 1997). We have built the development and feasibility of ward interventions in collaboration with our lead participating site to ensure we take an approach that can respond to the complexities of implementing change within healthcare systems (Holmes, et al, 2017).

Study Results

Resistance: a routine part of everyday care

We treat 'resistance' as an emergent concept and we found that it surfaced in the ways staff interpreted and responded to and assessed people's behaviours within the ward as legitimate or not. Staff were aware that there were patterns of resistance throughout their shifts, however, they were typically not able to recognize the potential triggers or patterns of resistance for individuals within their bay. Resistance was typically seen as a feature of the dementia diagnosis, and a behaviour to be expected from this patient group within every shift.

HCA discusses the ward: It goes in waves, it's not bad today, I am watching (bed number). As we are talking a nurse comes over and says she will take over the 121 work and sits in the chair and fills in the patient chart and takes a new 'side room checks' document for him [Site B day 3]

However, despite these routinely observable patterns or 'waves' across shifts, staff interventions in response to perceived resistance, could be highly variable.

Overall, we identified extremely high levels of resistance to care amongst people living with dementia within acute hospital wards. It was a common feature of ward life; every patient living with dementia we observed admitted within the acute hospital ward, resisted care during our period of observations. In total, we identified 1,052 (T&O=523, MAU=529) incidents or episodes of resistance to care. This could manifest in a number of ways (in order):

- Attempting to get out (and getting out) of bed, standing and walking around. This includes: wanting to go home, trying to leave the bay, ward or hospital; going to other rooms or areas of the ward or unit; pulling at the bed rail; approaching the nursing station.
- Verbal and physical cues. This includes: shouting, angry, sounding agitated; crying, screaming, sounding distressed; asking to go home, see family or be taken somewhere; biting, pushing, pulling, spitting at, or holding on to staff; refusing to leave the ward, be transferred, or discharged.
- Not submitting to the timetabled rounds of the ward. This includes: mealtimes, food and drink, medication, personal care, observation rounds and examinations.
- Removal of equipment. This includes: pulling or removing IV, gastric tubes, dressings, catheters, nebulisers, cannulas, oxygen mask, and other medical equipment crucial to care; pulling sheets off the bed or removing clothing.

Resistance, in whatever form it manifested, was always identified and interpreted by staff as a feature of the person's dementia diagnosis that signified their lack of capacity, and as such was to be overcome or managed. Similarly, it was often interpreted by family carers and visitors as 'bad behaviour', something to be embarrassed by and to apologise for. However, we identified that resistance was typically a response by people living with dementia to the everyday organization of their care within the wards (fixed timetabled routines), and the ways in which ward staff delivered care at the bedside. Importantly, these findings were represented in our data within all hospital sites, regardless of ward, time of day, day of week etc. Our ethnographic approach allowed the researchers (KF and AN) to spend extended time within and across shifts observing specific ward bays and the individual staff and patients within them. These extended periods of observation almost always revealed the potential underlying reasons or triggers for a person's resistance to aspects of their care or their admission. These were typically rational to that patient's present ontology and perceptions.

Ward staff responses to resistance was either to (1) ignore resistance, particularly if the behaviour was judged not to be an immediate risk for the patient or interfering with the organization of the ward; or (2) to intervene and manage, particularly if they believed that the provision of care was essential, that there was an issue of patient safety, or that it was interfering directly with or delaying the organization, timetables and wider work of the ward. The subsequent patterns of responses and interventions typically involved nurses and HCAs using multiple interactional approaches that combined language (orient to reality, instructions to be obeyed, the necessity of the institution, negotiation and bargaining) and the performative (a focus on completing essential care and containment of the patient at the bedside). These approaches created damaging cycles of stress for patients, families and for ward staff.

Physical Resistance: Getting out of bed, standing, walking

People attempting to get out of their bed or the bedside chair and, if that was achieved, to stand and walk, was interpreted by ward staff as a key form of resistance. This was the most overt and commonly observed resistive activity and was consistently high across all wards and sites (although slightly lower in sites C and D). This response to care was expressed by the most immobile patients, such as a person who was only able to clutch at or pull and 'rattle' the raised rails at the sides of their bed (for this patient group, if they were in bed, the side rails on the bed were typically raised); people sitting in their chair who were able to push down on the arms of the chair in an attempt to stand up; to people who were able (often unsteadily) to walk away from the bedside. However, this behaviour was only an immediate concern for staff and identified as particularly problematic if the person was able to stand and this was likely to lead to the person walking from the bedside and walking unsupervised within, or from, the ward.

In this example, an 86yr old woman who is living with vascular dementia has been admitted following a 'fall' and over one afternoon made multiple attempts to try to get out of bed. Every time she tried to leave the chair or the bed the HCA in the bay immediately responded to and repeatedly helped her to sit back in the chair ('let's sit you up for lunch') or lead her back into her bed. The team (particularly the HCAs) encouraged her to sleep, arranging the sheets and pillows and holding her hand to try to calm her. However, every time the patient appeared settled, she would immediately return to reach for something, to sit up, or get out of bed and stand. Her need to get up can be related to her immediate concerns and anxieties she expressed for her home, where her house keys were and who was picking her up and taking her home. Throughout this afternoon, staff responses to her started with distraction (lunch), to repeatedly questioning her 'what are you doing?', 'where do you want to go' and 'who?' and then escalated to giving her clear directives 'you need to rest' and enrolling the requirements of the wider institution by suggesting that other people having the power to decide whether she can leave 'We need to talk to the doctors first' and 'your sons will sort it out for you':

She is wearing her dark red cashmere jumper over her hospital gown and she tries to get out of bed. She has lifted the sheets off her legs and swings round to a sitting up position with her feet on the floor. She has been looking in her handbag and getting out of bed a number of times all morning and I think she was looking for her key and expecting to go home.. The HCA immediately goes over to her

Patient: ooh I am out of breath- she breath heavily

HCA: its lunch time, let's sit you up for lunch. She helps her to move slowly and gently from the bed to the bedside chair and puts lots of pillow behind her and tidies up the bed.

Patient: Thank you

The HCA tucks her into the chair with her mohair blanket over her legs with the trolley longways right in front of her, which keeps her in place. The trolley wheels are on the mohair blanket that trails on the floor and she stoops to pull it off- she is perilously close to the edge of the chair because there are lots of pillows behind her that also are pushing her forward. The HCA goes over from her chair at the other end of the bay to help her, puts the blanket on the bed and helps her into bed. [...]

Patient: It's a bit worrying they are supposed to come for me I only live over the way- she points out of the window next to her bed. She looks uncomfortable in bed and looking around for something on the bed and then gets her legs out of bed

The HCA goes over to her: what are you doing? She lifts her legs back up into bed and tucks her back in and moves the trolley next to the bed within reach. As soon as the HCA leaves, she leans over to the end of the bed with some difficulty and gets her mohair blanket and goes to get out of the bed and lifts her legs over the bed and her feet on the floor. The HCA returns: where do you want to go? Where do you want to go? She is kind and gentle in tone, but also sounds slightly exasperated

Patient: She's picking me up in a minute so she said

HCA: Who? The patient sits on the bed and shakes her head, she is trying to remember and the HCAs leaves and as she turns this patient tries to get up from the bed so the HCA returns and goes to help her to pull her hospital gown down, it has ridden up and helps her to move from the bed to the chair and sits her down and strokes her face- I am just over there- she points to the chair in the corner.

Patient: Where are you staying tonight?

HCA: I am going home

Patient: someone is going to pick me up, I only live across the way- she points outside of the window. The HCA holds her hand and strokes her gently and soothingly. I am waiting for someone to take me home

A young male HCA arrives and has joined them: take your medicine, eat well and you can go home! The HCA sits with her and a nurse comes over and relives her while she goes on her

break and helps this patient into the bed: you need rest. She puts the bed flat and helps her to settle down in the bed- she is curled up on her side and she places one pillow behind her back and one in front of her shins and covers her up. She tells her This is your blanket, and she makes sure the mohair blanket is over her and she can see. The nurse sits in the chair and holds her hand and strokes it gently

Patient: I wonder who has the key? the nurse strokes her head gently. The patient sits up: what do you suggest?

The nurse is next to her: Nothing, you need rest, its best to sleep you need rest [...] The nurse leaves and the patient then pulls the blanket aside and moves the pillows and swings her legs out of the bed and sits on the side of the bed. The HCA watches her and goes over: Where are you going?

Patient: Someone is going to take me home I hope

HCA: We need to talk to the doctors first

Patient: Ok [...] where are the keys for my home?

HCA: I am sure your son or daughter has them, you must stop worrying about it, try not to worry your sons will sort it out for you [Site E day 11]

Although the bay team talked to her throughout this afternoon, as they did this they also continued to subtly restrict her movement to the bed and the bedside. They start with tucking the blanket over her legs and placing trolley in front of her in the chair, and then use pillows around her body, cover her up with the sheets and sit next to her holding her hand.

Getting up, standing, trying to walk, walking or walking unaccompanied within the ward was always discouraged and problematized as a form of resistance for all patients. However, this could be subject to a variety of responses dependent on the assessment of the immediate risks to the individual patient. Patients who evidently did not have the strength to follow through and achieve their aim of getting out of bed, standing or walking, were typically not assessed by staff as an immediate concern. Here, a 94 year old man with a fractured hip and pneumonia, continues to rattle and pull at the raised bars at the sides of his bed. However, even though he does this forcefully and over an extended period during this shift, an indication that he appears to be distressed, the team continue their focus on the routines that must be completed and recorded around him - the observation and medication rounds - to the people in the other beds within the bay. They do not check on him directly; he is contained safely within the bed:

The HCA returns to the bay with the mobile Blood Pressure unit from taking the measurements from the patients in the cubicles, the nurse finds the yellow charts for him have returned and are open on the trolley at the end of the bed and updates them. He is a very thin man wearing a hospital gown lying in bed - the sides of the bed are up and he is holding on tight to the left side bar and shaking it and rattling it as if trying to get out. He moans and sounds very distressed. The nurse continues and moves on with the medication round to other patients in the bay and as she is doing this, he holds tightly onto the raised side bar on his bed, continuing to rattle and pull at it. [Site A day 4]

The organization of care and the routines within the ward meant that often staff were not able to see this behaviour as a priority for care, and if they did, the perceived priorities of the ward meant that they often appeared unable to respond. However, walking or walking unaccompanied within the ward was almost always discouraged and problematized as a form of resistance for all older patients. Staff typically questioned older patients who were walking and the language discussing walking was as a danger or a form of 'escape', often using humour to reduce this behaviour:

The HCA wakes a patient up for her observation and medication. As she does this a younger man in his 20's from the bay at the end of the ward walks past, he is just wearing his underpants and is holding a wash bag and pushing his mobile drip heading to the bathroom. At the same time an 87 year old woman with a diagnosis of dementia who has been admitted with a fracture is trying to get up from her chair. She is wearing a full length quilted dressing gown that is buttoned all the way up to her neck (it is the middle of summer and very hot in the ward). It is purple with tiny sprigs of flowers like heather all over it. As she does this, the HCA calls to her: where are you off to! She responds that she is heading to the toilet and the HCA is relieved: oh I thought you were going to escape! She helps her to reach her walking frame and stands behind her as she mutters 'I know I know I know....' All the way to the bathroom and back. Afterwards she turns to the HCAs and smiles and says 'thank you so much'. The HCA gives me a look as the young man from the bay at the end of the ward walks back from the bathroom in just his underpants. [Site A day 1]

Throughout, judgements were also made about the person, their condition and whether this behaviour is legitimate and to be permitted within the ward. A man in his 20's was able to walk up and down the ward in his underpants without comment, however, this woman who has a diagnosis of dementia is met with close control over her movements.

Verbal Resistance: Shouting, anger and agitation

It was common for people who were shouting, agitated, or angry in the ward or bay to receive the most attention from staff. This was a key response across four of the sites, and was only low in site C. This behaviour often resulted in the individual being assigned one-to-one agency care ('specialing' a patient means that their needs have been assessed as so great that they require care at all times and they will be assigned a private agency HCA to provide one to one care) within the shift. The key responses from ward staff to this behaviour was to (in turn) orient people to where they were, direct them to calm down, and to rationalize with them. Throughout these processes there were typically multiple attempts to contain the person at the bedside, which could lead to them being supervised more closely, which in turn could cause the patient further anxiety, and trigger the ward procedures of 'specialing', a DoLs order, and one-to-one HCA care. Here, a 77yr old man with a diagnosis of dementia, has been in this ward for over 5 months, and although he is 'medically fit to leave' there is no package of care organized for him. He is in an individual room and is 'specialed' with one-to-one HCA care and although he has difficulty communicating, he is clearly waiting for his lunch. Shouting 'HEY' is his main form of verbal communication during the day and as he waits, he gets louder. The one-to-one HCA and the team passing join in and they repeatedly switch between encourage his excitement about the arrival of lunch and then use a number of techniques in an attempt to calm him down when lunch does not arrive on schedule. They orient him to where he is 'you are in your room', direct him to calm down, and then rationalize with him 'what do you want?'

I am walking through the corridor with 2 HCAs and the person in the side room shouts for us. The door to his single room is open and he is sitting in the chair looking out wearing hospital pyjamas and red hospital socks and he has a large knitted hat on his head. The one to one HCA is sitting very close and opposite him filling in his bedside chart. We go over to see him and he holds up three of his fingers to indicate that there are 3 of us. The HCA with me asks him what he is having (pasty) and asks him what sauce he would like to go with it and she gets him sachets of sauce to have with his chips: red, brown, and vinegar. He is now very excited about the chips. I introduce myself again and we talk about the hospital food and he tells me the food is terrible and then covers his mouth like he shouldn't say this and we laugh together and sympathise. He gets more and more impatient for lunch- Hey HEY and asking for the time when the chips should arrive and the team count down the time -

15...10....6.... Patient: HEY HEY Give it, give it, give it me. He is now getting very angry and

agitated and repeatedly shouts HEY HEY as people pass his door

HCA: What's the matter now? listen you need to calm down

Patient: Hey hey

The HCA continues to talk to him and repeatedly tries to calm him down [...]

Patient: HEY HEY HEY

HCA: She jokes and laughs with him and is very positive- Your pasty is coming and ice

cream
Patient: HEY

HCA: Are you alright? Patient: NO I AM NOT

HCA: What do you want? Lunch fails to arrive and the team call catering. When it finally arrives the pasty is cold, the chips are soggy and his agitation, unhappiness and frustration

increases. [Site D day 5a]

This behaviour suggested that the person was attempting to communicate with ward staff. Importantly, it was also a response to the ways in which care was organized and was being delivered at the bedside, especially (as in this case) if this does not meet the patients expectations. In the case above, staff encouraged the patient's excitement about his lunch only for it to fail to arrive on time, and be unappetising and inedible when it did.

Resistance towards the timetabled rounds of the ward

A common way in which ward staff identified resistance to care, was when patients failed to submit to the timetables of the ward. This included mealtimes, food and drink, medication, personal care, observation rounds and examinations. This study has a specific focus in examining mealtimes and medication rounds and resistance to food, drink and medications. Thus, we examine these aspects of routine care in more detail within the wider context of the timetabled rounds of the ward, which includes: mealtimes, food and drink, medication, personal care, observation rounds and examinations.

Mealtimes, food and drink:

The delivery of food and mealtimes was an aspect of the routine timetabled care within wards that typically prompted resistance to care (this was consistent across sites). Mealtimes was also a point in the timetable where there was an emphasis on speed and efficiency, particularly for people who were assessed as needing support to eat. This was often in the form of a clear communication from a person that they do not want to eat (verbal or physical) or the absence of communication (silence in response to often repeated requests). In response, this was almost always viewed by staff as a feature of the person's dementia diagnosis, a sign that they did not have capacity and a form of resistance that must be overcome, emphasising the importance of food and nutrition to recovery. Even when people clearly indicated that they did not want any more food, staff would typically make a further attempt to encourage them to eat more before stopping. However, staff responses to individual patients was highly variable and while for some patients there appeared to be more flexibility that allowed for a wider range of behaviours and responses around mealtimes and eating, for others, mealtimes became a battleground. This meant that mealtimes often triggered anxiety and the potential for this resistance to escalate quickly into the person refusing all care.

Needing help with mealtimes seen as a feature of the condition

For ward staff, a person with a dementia diagnosis typically required help with eating meals and not being able to eat independently was viewed as a key feature of the condition. In addition, rather

than recognising that the issues around food and food refusal may be signs of other underlying issues (confusion, anxiety, unsuitability), they are expected as routine and inevitable aspect of an admission for this population. This was also seen as a feature of their condition, which was either permanent or could become better managed once the patient was discharged or transferred to a specialist setting. The issues around food were recognised as something that manifested within non-specialist acute wards and short-stay units like assessment units, but could not be prevented or solved within them.

Even when a person clearly refused food either by keeping their mouth closed, by pushing food away or by stating that they do not want food, staff typically continued to try to encourage, persuade or put food into a person's mouth. This could be obviously distressing for the person, although this impact was typically not recognized by ward staff. In these interactions, the meal is no longer pleasurable and instead becomes a confrontation between the member of ward staff and the patient. The potential for this resistance to escalate quickly into the person refusing all care was high during mealtime interactions. When resistance and refusal was exhibited it could inform staff perceptions of that person and their understandings of the impact of their dementia diagnosis and their capacity to make decisions, and create anxiety and confusion for the often-admonished patient. Here, despite finding speech very difficult, this 96 year old woman with a diagnosis of dementia who has had surgery for a fractured hip is able to clearly and angrily articulate that she does not want any lunch. However, this is interpreted as a feature of her dementia to be overcome and the volunteer continues to place a forkful of pureed food in front of this person's closed mouth, starting with encouragement 'just a little bit' but quickly moves on to a directive 'you've got to eat', 'open up'. This is seen as essential work and this mealtime volunteer continues this approach and only stops once this patient has accepted a tiny bit of dessert into her mouth, even though she then spits it out. The volunteer does not wipe away the food around her mouth and leaves the bedside, clearly frustrated at her lack of progress. The experience of the repeated attempts to spoon puree into her mouth despite her saying no, appears to leave this patient in an increasingly anxious and fearful state and she resists all further care that day:

This woman is lying in bed and I haven't seen her leave the bed yet. She is wide eyed looking around the bay and has been talking quietly all morning. The mealtime volunteer arrives (a tiny woman wearing a tabard and who appears to be in her 70s); she places the tray of lunch on the trolley and sits on the chair next to her and then leans over the rails to talk to her. The plate has pureed food- a third is bright orange, a third is white and a third is a dark grey with bits in (I think this is pureed stew). She uses a large metal fork and puts pureed food on it, it is orange but hard to make out what it might actually be, possibly carrots? She puts the fork to her closed lips: Just a little bit Patient: I don't want any. This is the first time in two days I have heard her say something that is clear and easy to understand- it has clearly taken her a great effort to articulate what she wants and she sounds very angry. However, the volunteer continues in an upbeat and encouraging tone: You've got to eat, just a little bit. She keeps the fork with the pureed food hovering at her mouth: just a little bit! She leans further over the side rails of the bed and holds the fork close to the patient's mouth. I am standing next to the sink and bathroom in the corridor opposite the bay and the volunteer turns and talks to me: I am a volunteer, I can't force them, I can only try. She puts the fork back on the plate and moves on to the dessert. She takes a large metal spoon with a similar looking yellow/orange puree on it and holds it in front of her mouth: just try a tiny spoon, listen, just try, unless you try you don't know, just try a little bit, open up just a bit. She does open her mouth a bit and the volunteer spoons a tiny bit into her mouth. But she looks extremely unhappy at this and uses her tongue to push the puree out so that it sits on her chin at the side of her mouth. The volunteer stops trying at this point and for the rest of the morning the yellow-orange puree sits on the side of her mouth. As the volunteer puts the tray away and gets ready to

leave the bedside she turns to me: I had to give up, she was the same at breakfast...I told my son if I get like this take me to the vet! She leaves and the nurse arrives with a syringe of liquid medication to take orally, but she won't take it and the nurse leaves and moves on to the next person. This woman is now talking in a low voice to herself, fiddling with her waistband (I wonder if this is anxiety or she needs to go to the bathroom) and looks around the room. She now looks very, very, anxious [Site E day 2]

An extremely common approach was for ward staff to emphasize nutrition and the importance of eating to the patient as they continued to try to put food into their mouths. Here, the team discuss the patients in their care and who needs 'feeding' in the bay (people living with dementia were often referred to not as people needing mealtime assistance, but given the label 'feeders' who needing 'feeding'). The nurse ignores this person's concerns even though this 95 year old man who has had a hip operation following a fall and has a diagnosis of dementia repeats a number of times that this is 'the worst day of my life', the nurse continues to encourage him to eat, emphasizing the importance of nutrition. Although he clearly expresses that he does not want help, she takes the cutlery and puts a spoon of food directly into his mouth. This encounter is also punctuated by his chair alarm which is repeatedly activated by his movement in the chair.

Nurse: hello, have you had lunch

Patient: everything in the world has happened to me today all the worst things. He holds her hand and she leans low over to talk to him: will you have some lunch for me, were having a party

Patient: I won't be able to eat it all

Nurse: your dinner is here my darling she speaks quietly to the HCA are you ok to feed him? She turns to him: if you can eat ice cream you can eat some actual dinner, you eat too many sweet things you need some nutrition. However, he is so shaky that by the time it gets to his mouth it has fallen off the form and is empty and in addition, his seat alarm keeps going off. He tells them: I am sorry I can't eat it

The team offer help but he rejects this: I don't want help and the nurse pops a spoonful in his mouth and he pulls away: Sorry love it's not very healthy to eat chocolate. He responds: I can honestly say it is the worst day of my life, I won't have this (main course) but I will eat that (ice cream) and he gets the ice cream and take bits of it with his fork as his chair alarm goes off intermittently blaring loudly throughout the meal. [Site C day 7]

The impact of repeated attempts to encourage people to eat and the approaches staff used to try to 'feed' someone was that it typically made people angry and increased their anxiety and distrust of staff. Here, the HCA's encouragement becomes increasingly forceful and despite appearing to ask this patient if she wants to try it, and despite the patient pushing her hand away, continues by repeatedly putting large spoons of food near her mouth. It is only when this patient, who is a tiny woman confined to the bed (the 96 year old woman with a diagnosis of dementia who has had surgery for a fractured hip above), makes another attempt to communicate her wishes by forcefully pushing her hand away that she stops. This is a surprisingly fast movement, which indicates her underlying anger and frustration. The HCA is shocked and is shaken by this and the patient looks increasingly anxious and fearful during the rest of the shift.

Lunch arrives and there is a strong smell of cooked food in the ward. The HCA takes this woman a meal of pureed sausage, pureed peas, mash and gravy: I have a lovely lunch here for you, do you want to try it? She is sitting next to her and puts a clean linen pillowcase across her chest and gives her a large metal spoon: you try it, hold the spoon. She helps her told the spoon and also holds it and guides it into the mash and then guides it towards her mouth. But this woman pushes her hand away from her mouth and puts the spoon back onto the plate. The HCA then once again uses the spoon to scoop quite a large amount of

mash and gravy onto the spoon and guides it back to her mouth and she takes a tiny bit. The HCA is very encouraging and moves on to the dessert- I have apple and custard (it is in a tiny pot). She sits next to her and puts a bit on a large metal spoon: Let's try again, she tries to put the spoon into her hand. In response, this woman takes it and pushes it away extremely forcefully, it is so fast and unexpected that the HCA jumps out of the chair in shock. I ask her if she is ok and she responds- You never know! And she turns to say to the woman: let's stop there. However, this patient now looks very anxious and wide eyed as she sits in the chair [Site E day 5]

The frustrations staff experienced around mealtimes could quickly become visible and they sometimes became angry and exasperated and raised their voice to patients. Although patients not eating a meal was seen as resistance to be overcome, for staff this also seemed to signify their own personal failure in delivering an essential aspect of patient care. Here, the HCA repeatedly tries to encourage this 82 year old man with a diagnosis of dementia to eat lunch:

The HCA encourages this patient to have dinner: have your rice pudding, how do you know if you don't try it? She raises her voice slightly and sounds exasperated: wont you try it! No? ok that's fine but it's a shame. This man is lying in bed only slightly propped up in hospital pyjamas and as she writes in his bedside notes she asks again: Do you want something to eat? Shall we try again? Don't you like sweet things? Do you want a drink before I go? He responds: No [Site B day 13]

Interrupting, taking over and hurrying up can lead to refusal

Mealtimes was also a point in the ward timetable where there was an emphasis on speed and efficiency, particularly for people who were assessed as needing support to eat. Here, this 95 year old man who has been admitted with a hip fracture following a fall and has a diagnosis of dementia is eating his meal without support, although he is doing this slowly. The HCA joins him and immediately interrupts, encourages him and takes the fork from him and tries to 'feed' him a forkful of food. In response, he refuses any more food, shakes his head and crosses his arms defensively.

The HCA goes over to this man, she usually 'feeds' him, but when she goes over to him, she brings a chair and realises that he is eating by himself. She puts the chair next to him and sits very close to him: do you want some gravy with that? He is taking tiny bits of food onto his fork and eating it. She leans in- you are going to do some exercises (physiotherapy) so you need to eat to get your strength up

Patient: That's a good idea

HCA: One more forkful? He shakes her head, but she takes the spoon and tries and gives him one more spoonful: if you have one more spoon you will have eaten half your dinner. He shakes his head and crosses his arms- he looks uncomfortable. The HCA asks him: Are you ok?

Patient: I've got the bloody itch (itchy back he has been complaining off on and off all week)

HCA: You don't want to eat more?

Patient: No. [Site C day 6]

Even when people clearly indicated that they did not want any more to eat, ward staff typically would make a further attempt to encourage them to eat before stopping.

Mealtimes often triggered anxiety and a very common source of anxiety stemmed from people being worried that they did not have money to pay for their meal or drinks. It is not uncommon to see patients begin searching for wallets or handbags (that they often do not have, which created a further source of anxiety and distress) as meals were served (this is particularly the case in MAU's

across sites). This could lead to real distress, and we observed several patients breaking down in tears at the thought they would be unable to pay for their food. While this was an issue that was often resolved by the presence of an attentive member of staff reassuring the patient, it could take more covert forms that go unnoticed. Here, a 96 year old man with dementia and 'increased confusion' and admitted following a fall responds to the tea round:

The team trolley comes into the bay and the HCA asks him: fancy a cup of tea? He comes back over to us and asks: is there anywhere I can get a coffee? In response, I offer to make him one and he starts fishing for money in his pockets. The HCA and I both say: you don't have to pay for it, coffee and 1 sugar? The HCA takes him a coffee (in a red plastic mug) and says: here you are duck. As she does this, the man in the bed opposite struggles with the biscuit wrapping and asks her 'do I have to pay for this?' She replies 'No its free darling' [Site B day 6b]

Supportive and enabling approaches to mealtimes

Importantly, mealtimes could be very supportive and enabling routines within the ward timetable, with staff taking time to support, encourage and help people to eat. Mealtimes could also be a quiet time in the bay; a point of communication and this could be transformative for the person living with dementia. Once people started to eat, this could have an immediate visible impact on the person and it could be a time when people would 'come back into the room' (ward sister site C) and their alertness and ability to communicate would visibly improve. Here, the ward housekeeper is with a 79 year old man with 'acute confusion', she starts by fully supporting him to eat breakfast and as she stands over him at the bedside, repeats a number of times the instruction to open his mouth. He finishes the porridge. She then asks what he wants, how he likes his tea and encourages him to help himself and hold the cup and he drinks it all. This is transformative and he becomes able to communicate with her, he is alert, his eyes focus on the ward around him and he starts to show an awareness of the room and his surroundings.

The housekeeper takes the bowl of porridge (it must be cold by now) and stands over him at the bedside. She uses a plastic teaspoon and takes some porridge: Open your mouth....open your mouth....open it....Just one more 2...open your mouth....wide....wide....thats it...porridge....He has finished the porridge and this is the first time I have seen him eat. She puts the bowl on the tray and turns to him: Shall I make you a cup of tea? How many sugars? He responds to her- again this is the first time I have seen him communicate. Later, she comes over to me where the tea trolley is and she talks about him: I know the tricks of the trade! I am not just a pretty face! I know how to do this! Don't use a metal spoon, use a plastic spoon, it is gentle and bends in their mouth and it helps them to eat, he had all his porridge. I daren't try the yogurt yet, but I will try him with a cup of milky tea. She makes a very weak milky tea in a sip cup and adds a straw and says to him: I have a nice cup of tea for you, she takes it to him and then comes back and adds some more milk. She returns and stands over him holding the cup and he drinks a bit through the straw. As she leaves he is sitting in bed with his eyes open and looking around the room. [site B day 4]

Breakfast was a particularly powerful routine at the bedside when staff were able to take time to support people. This could make a difference to improving the person's alertness. Mealtimes could be a time for communication. This work included staff asking the person about their family and also sharing personal information about their lives. This emphasised the importance emotionally of food and the potential for mealtimes as a time of connection and communication.

Drink and hydration

We did not observe people living with dementia resisting drink or opportunities for hydration. In contrast, we observed high rates of patients without a drink within reach, unable to drink from the cups they were provided with, or their requests for a drink nor prioritized by staff. In only one ward (Site C) did we observe that it was regular practice to focus on optimizing hydration and to consistently offer people a drink of water or tea during the majority of the routine encounters at the bedside. Within this ward there was a regular emphasis by all staff on supporting hydration at the bedside during every encounter with a person. Here, 'sippy cups' were used, but they also used the simple straw in the spout in addition to this so that people did not need to tip the cup to drink from it. Staff handed people the cup to hold and emphasized the importance of drinking. Here a 95 year old man with a diagnosis of dementia has been admitted with a hip fracture:

He is sitting in his own pyjamas with the tv monitor low and in front of his face, his glasses are on and he is watching the news he dozes and watches a bit and then dozes again. He has a sippy cup of tea in front of him and a glass of water with a tiny bit of water in and a straw. The HCA comes over: can I do your blood pressure? She places the cuff on his arm and the BP machine beeps: there you are all done, its fine, you just need to keep drinking water for me darling, I want that whole cup done! She passes him a glass of water with a straw and he drinks it [Site C day 7]

However, overwhelmingly, within all other wards, it was extremely common for glasses of water at the bedside to be empty, or to have low levels of water in them, or to be out of reach. Across the wards, as well as traditional institutional style china or ceramic cups, saucers and mugs, 'sippy cups', tinted and coloured plastic water glasses, straws and thickener were all used to provide water, tea and coffee and to support hydration. However, these technologies were not always enabling and could be difficult or confusing for people to use. The 'sippy cup', a spill-proof drinking cup originally designed for toddlers, was commonly used for people living with dementia in all wards. Here a 94 year old man with 'memory problems' is trying to drink from a plastic sippy cup. However, to drink from it, it must be raised high and tipped up to position it at an angle to drink from. Although he tries a number of times, he does not have the strength in his arms and cannot raise it high enough.

He is lying propped up in bed, the sides of the bed are raised and on the right side the side bar is padded with a thin mattress and the trolley is over him in the bed with a sippy cup of water. He keeps his eyes closed and takes the sippy cup and puts it to his lips he tries a number of times to suck water into his mouth, but can't lift it high enough to get any water in his mouth and he puts it back on his tray [Site B day 6b]

The use of sippy cups was common and although it is also a technology designed for someone with a weaker grip or someone lacking dexterity, its use did not take into account the strength needed to lift it high enough to use it effectively. Importantly, for people living with dementia, even if water was within reach, it may be that they either do not recognize the glass, mug or sippy cup in front of them or recognize that they are thirsty. This underlying dehydration only became apparent during the course of a shift, when they were passed the glass of water and in response would often hungrily drink it all.

Resistance to medicines and the medication round

The majority of encounters with patients during medication rounds were unproblematic. However, when examined across whole bays, wards, or units, this was a common occurrence during each medication round and it was rare for a medication round within a bay of 6-8 people to be completed without any resistance and it was usual for the routine to stall with a person at one or more of the bedsides (with particularly high rates in sites A and C). Routine cycles of conflict and resistance

during the medication round was observed that were repeated every shift, often with different staff, rehearsing and duplicating the interactional performance and routines of the medication round during encounters with the same patient. This often escalated into conflict and led to the patient being labelled as 'difficult'. Resistance to medication was not an exclusive feature to patients living with a dementia or with cognitive impairment, although they were more pronounced within this group, this was common amongst older patients within the wards.

Medication rounds were a key ward routine where staff expected that patients would exhibit resistance. The medication round was always a time of increased urgency and often increased anxiety for nursing staff, which was driven by perceived constraints within the ward timetable in completing this task and the importance of patients taking all of their (typically multiple) medication. Staff expressed a clear sense of relief and accomplishment if the round was completed without perceived resistance and delay. Common reasons for people resisting, guerying, or challenging their medication included that the medication offered was not the same as their home prescription (including variation in brand, dose or delivery of familiar medications) or apprehension of medication that they feared would have side effects such as nausea. Although this was a typical feature of the routines, this is always viewed by staff as problematic and resistive, and appeared to increase their anxiety, stress and frustration. Here, the nurse focusses on providing repeated information on who has prescribed this (the 'doctor' and the 'psychiatrist') new tablet. It is only then this patient, a 70 year old woman who has been diagnosed variously with 'dementia', 'delirium on a background of dementia' and 'manic/hypermanic behaviour' demonstrates in a number of ways that she would not take the tablet by disposing of it in her teacup, followed by 'no', that this nurse gives up. She writes this in the notes, but predicts that she will be criticized for not completing this.

The nurse goes over to her with medication in a pot. She has a loud voice and is very strident: I have one little tablet from the doctors who saw you today. She puts it into her palm and the woman looks at is closely: It's not the blue one. She picks it up and puts it in her tea cup straight away. The nurse is clearly extremely exasperated: can I explain to you! Because you have been seen by the psychiatrist today he has given you this. Patient: No its rubbish. She is very clear that she doesn't want it. Nurse: He has prescribed it, Ok you don't want to take it. She is very exasperated and puts the rest of the medication into her personal drug cabinet, saying to herself and the wider ward: It's the first one that has been prescribed! She sounds very frustrated and writes in her bedside notes and says to me: They will say you haven't tried! (it's the lorazepam)[Site E day 8]

This frustration is also linked to the visibility of the medication round. When a medication round has not been completed, nursing staff appeared to feel exposed to the scrutiny of others in the institution for this apparent failure to complete the task. However, the medication round and the persistence of the team trying a number of approaches, despite the person clearly stating 'no', could cause anxiety that has repercussions for the person's care throughout the shift. As for this 96 year old woman admitted with a fractured hip, her anxiety increases and she refuses all further care during this shift.

The HCA goes over to her: I will just wipe your mouth

Patient: Why? HCA: You have food around it. However, she doesn't like this and makes it clear and as this is happening, the nurse arrives with her medication in a pill pot- two pills and the HCA takes one and tries to help her take it by putting it in her mouth: can we give you this tablet? can I pop it into your mouth and a bit of a drink? Patient: No. HCA: With the dessert? She seems to agree. Do you like custard? The HCA puts custard on the spoon and places the tablet on top: Try a little bit. She puts it close to her mouth. Patient: No. However, the HCA continues to try to get her to take the tablet: this medicine will help you! The tablet is now just in her mouth with a bit of the custard, but she spits it out and it is in

her hand and she tells the HCA: you are a bloody nuisance you are. The HCA picks it from her hand and puts it in a tissue and puts it in the bin. However, the HCA perseveres and returns with another tablet, but does not manage to get the tablets into her mouth and in the end places the second tablet also covered in custard in a tissue and in the bin. The HCA leaves, but now this woman is quite distressed and talking to herself and looks very anxious and wide eyed. Some time later, the nurse does a quick sweep clearing up the bay and goes over to this woman who is now talking to herself and looking anxious. She checks the pill pot, sees that it is empty and puts it in the bin, she clearly assumes she has taken the tablets [Site E day 4]

For many people, resistance and rejection of their medications was part of their wider rejection of all care.

Attributing agency

A key element to conclude most nurses' medication routine at the bedside was to check that the medication had been swallowed, usually by asking them to open their mouth to check that it has been swallowed. However, it was still common for patients to spit them out, hide or to throw them away. There was sometimes a recognition that for some patients this was a reasonable response 'I can't blame you'. However, ward staff typically interpreted this as a wilful and resistant act by the patient, who were described as 'naughty' or being 'crafty' and 'he's a monkey', as in the case of this 95 year old man admitted with a fractured hip:

This man tells the nurse that his pain is 'nasty' The nurse responds: here are some painkillers, if you eat more I can give you stronger ones, here you are, hold it. She gives it to him. The medic comes over to the bedside and interrupts, emphasising to this patient: the nurse has given you this to help you. However, he wants to save it for later and so the nurse encourages him to take it now: do you want some more dinner? do you want to try the tablets now? the painkillers are for the pain. In response, he puts it in his mouth, puts the next one in and he sips his drink. He spits it out and the student nurse notices and repeats: It's a painkiller so you don't have pain, you remember you broke this leg, you need painkillers, if you need more I can give you more, we just don't want to be in any pain. In response, he has quietly spat them out and there are large pieces of tablets surrounding hum on the floor. The student nurse tells me that he showed me his mouth to show me it was empty and then spat them out and told me to keep quiet! She giggles and cleans up his tablets from the floor and shows him: naughty!

He demonstrates how he spat them out undetected: I just went like this! [Site C day 5]

Performance – putting tablets directly into the mouth

The most common and widely used approach across all sites to deliver medication to people living with dementia was to put tablets directly into a person's mouth. This technique included putting tablets on spoons, putting spoons or tipping pill pots into a patient's mouth continuing during these encounters. Tablets were also crushed or mixed in with food, particularly desserts such as yogurt, mousse or trifles. During the work of completing the medication round sometimes the nurse would tell the patient they were going to place tablets in their mouths, with the patient opening their mouth to receive them. Here the nurse is with a 73 year old woman who has been admitted following a fall and described as 'pleasantly confused'. She describes the medication and that she is going to place them in her mouth and she continues to talk to her and explains what is happening step-by-step throughout this encounter:

Nurse: we have your tablets, can you open your eyes for me? I have two tiny tablets here I am going to put them in your mouth. If you can swallow them for me, one at a time, I am going to put it in, well done darling. I am just going to pop this one in now. They are a bit big aren't they, come on. Her tone is very gently and relaxed, no sense of hurry. Well done darling, it looks like we have some more tablets! Sorry.

Patient: I keep thinking we are done but she gives me more!

Nurse: just one more tablet, just one more, she touches her shoulder and leans over her and puts the tablet in her mouth and helps her with a sip of juice [Site D day 2a]

However, this technique was used extensively in response to a patient's resistance to taking them, using spoons to open often unresponsive mouths or to tip them into a mouth that did not seem easily able to respond and swallow them. Here the nurse expresses her frustration and annoyance that the patient, an 86 year old man with a diagnosis of dementia who has been admitted with pneumonia, has spat the tablets out. In response, she uses the teaspoon with pills on to pry open his mouth and tries to push them back into his mouth.

Nurse: Oh you have spat it out! It's me again! It's a pain in the backside, you need to take them! She uses a plastic teaspoon to push the tablets into his mouth: shall we try one more? She puts the tablet on a teaspoon again and puts it in his mouth and tries to give him a sip of juice: Can you feel it, it's your cup? She puts it in his mouth: A bit higher darling, she helps him and guides it up to his mouth and he takes the cup and raises it to his mouth. She sits next to him: have another sip, have you got it? have a sip darling. She guides the cup to his lips [Site B day 4]

Often this was part of an extended series of negotiation and pleading with the person. Here, the nurse kneels in front of the person and after describing the tablet and asking her to put it in her mouth, he pleads with her and seems exasperated and at a loss as to what to do next. The HCA sitting with the patient, who is a 90 year old woman admitted with a fractured hip and has a diagnosis of Alzheimer's, takes the tablet, puts in her mouth, and follows this with a spoonful of ice-cream.

She is sitting at the dining table in the middle of the room. The nurse comes over and kneels next to her: look at this, here is a tiny tablet, put it in your mouth. In response, she picks it up and throws it down. Nurse: try my little tablet! he is now pleading with her and is slightly exasperated, but it also betrays his tiredness and that is not sure what to do next. The HCA comes over, takes the tablet, puts it in her mouth and scrapes the last of the ice-cream onto the spoon for her to eat [Site D day 11]

This emphasizes the challenges for staff as they try to negotiate with people living with dementia. For staff, delivering each patient's medication was viewed as what should be a short interaction, a fast-paced task within their overall routine and the ward's timetable. Although the medication round could often start in a relaxed way, as it progressed, it often appeared to take longer than their timetable allowed, which led to nursing staff often becoming anxious and feeling they were taking too long and needed to speed up the process at the bedside to complete the round. It was typical for nursing staff to appear frustrated at the length of time the medication round was taking. Even within this round, where there were little distractions and the bay was quiet and calm, this nurse sighs and believes it is taking her too long.

Overall it is very quiet in the bay, just the low hum of the bed pumps and is with the first person in the bay to complete the medication round. It takes time for this first patient, who is with a person living with Lewy Bodies dementia and has been admitted with a fractured ankle. She hands him all his tablets and he picks up the pill cup and shakily tips them into his

mouth and then takes some water and drinks about an inch of water. There is also another soluble medication in a cup and he takes it a bit at a time. The nurse puts the packet of medicine back in the box and says to me and the wider ward: the medication round, it's very slow! [Site B day 5]

Trigger for patient anxiety

During the medication round, nurses were very task focused, working on dispensing medication at the bedside. However, this focus on completing the task meant that other features of the interaction or the impact of the round on the patient at the bedside became invisible. As in this case, this person seems agitated (he is 94 years old and admitted with a fractured hip and pneumonia, and has been agitated for some time and pulled out his IV port previously) and although she reassures him, once he has taken the tablets, she does not enquire further or checks if there is an underlying reason for this agitation. She continues with the medication round and moves on to the next patient.

The HCA is with him and she talks to him as she works: I have some blankets for you, you seem a bit cold this morning. She also wipes up the floor after the spill from the IV unit that he had pulled out and uses antiseptic wipes to clean the IV unit. She keeps gently talking to him and he responds: I SHOULD BE IN THE CLINK. He is very fidgety and this increases. The nurse comes over to him: you're confused this morning, you are not well at the moment, you've got a chest infection, and she repeats this in a calm and clear voice and moves on to the next bedside. [Site A day 4]

The medication encounter appears to have left this person additionally distressed and the HCAs is left to follow-up on his care at the bedside. She reassures him and focusses on orienting him to the reality of his situation.

Personal care

Ward staff always explicitly sought the person's permission to carry out the personal and intimate care of washing, changing clothes, changing continence pads, changing sheets. However, they also typically started work on the body immediately and continued with the task at hand (whether the patient assented or not), highlighting the perceived essential nature of this care, should the patient not accept or submit to their request. In this context, the person often had difficulties communicating verbally, was often woken up by the request, and appeared anxious or afraid of moving, and so any negative response they had to this type of request was often in the form of a physical act of resistance (pushing away, pinching, hitting out) or verbal (shouting, screaming, telling staff to stop), while staff continued with their focus on completing care. This was the case across wards and sites, although resistance to personal care was lower within site E. For staff, this work was of additional importance to the wider ward of presenting a neat, and tidy patient, bedside and bay to meet the timetable, routines, and expectations of the ward:

The bay team are with an 84 year old man admitted with 'confusion'. They are behind the curtain and they chat to him asking him about his life: what was your job when you were young ,where did you work? Do you remember? In response, he whimpers loudly and they apologise if they are hurting him and tell him what they are doing: sorry it is hurting you, we will dry it now we will put some cream on. He cries out: OW, OW, AAAH, and this turns into a high pitched scream. The team continue to explain to him what they are doing: It's to protect you, some lotion for your feet ok? I will roll you over. Ok, we will roll you this side, hold onto the bar, well done, nearly there. We are trying to clean your back, well done! In response, he whimpers, and lets out a very high pitched cry [Site B day 5]

This was a routine feature of care that occurred 'behind the screen' and so the analysis is based on observations blocked by the privacy curtain or screen, but where verbal cues (the conversations between staff and to the patient) and physical cues could be picked up (by moving curtains, sounds and odours).

Observation round

Observation rounds were interpreted by ward staff as an essential feature of the organisation and timetable of the ward shifts and these observations (blood pressure, temperature, oxygen saturation, heart rate were typically recorded) must be completed and recorded within the patient bedside records (typically secured to the end rail of the bed). However, for people who appeared to have tiny arms, or very 'thin', fragile or bruised skin, the observation round and specifically the blood pressure cuff seemed to cause a lot of distress. However, any resistance to the cuff on their arm was typically interpreted by staff as a feature of their dementia, rather than a potential physical impact of the cuff, which was rarely taken into account (this was consistent across wards and sites). When faced with resistance, although in some cases staff stopped, moved on to return later, staff typically continued to attempt to take a recording, and as with the medication round (although less common), the observation round typically stalled at the bedside of a patient who resisted the BP monitor.

The HCA is carrying out the observation round and starts at the bedside of a 96 year old woman with a fractured hip and a diagnosis of dementia. She tries to put the BP cuff on her arm: just relax its nearly done. However, she is getting increasingly agitated, pulls at the cuff, is making moaning sounds and is clearly distressed and in pain from the cuff on her tiny arm. The HCA tries to reassure her: I know, I know, its nearly done, and she holds her hand in an attempt to calm her. But she is getting even more distressed: You need to relax, otherwise it won't read, I know it's not nice, I know, just hold my hand, just relax. As this is happening, a mealtime volunteer arrives with her lunch (a plate of pureed food) and sits down on the chair next to her and tries to distract her with the food: 'you've got your lunch here', as the HCA continues trying attach the mobile monitor to her arm and to get a reading. This continues for some time and the HCA stays with her and tries to get a reading and gives her further instructions: keep your arm still, nearly done, it's alright, just keep your arm still nearly done, it's alright, just keep your arm still for a minute, keep your arm still and eventually the HCA decides that this is not going to work and she is getting more and more distressed and is now whimpering. Throughout this, the HCA has been holding her hand and she takes the cuff off her arm. She continues to whimper, is very distressed and immediately covers up the arm where the cuff had been with a sheet to protect is and is looking around her and looks very anxious and afraid. [Site E day 2]

This cycle continued with this patient during every observation round that followed and had a detrimental impact on this person, who became increasingly fearful of ward staff. It was also had a wider impact on other patients in the bay who could hear her cries and also became distressed. However, for ward staff, this was viewed as essential work to be completed and recorded.

Removal of equipment: Treatments and Tubes

It was common for people living with dementia to try to remove physical interventions and treatments that were attached to their bodies. This included pulling or removing IV ports and lines, gastric tubes, dressings, catheter tubes, nebulisers, cannulas, oxygen masks, and other medical equipment. These interventions were seen by ward staff as critical to care and resistance was typically responded to by repeatedly tightening or securing the equipment in place or out of reach.

All is quiet in the bay and this 83 year old man admitted with pneumonia, hasn't moved for ages. His eyes are open and he is intermittently pulling at the IV port in his arm and around the bandaged area. It looks like a whole role of bandages have been wound round his entire arm to secure this in place and he is starting to cry out. He keeps tugging at the IV port and brings his hand and arm close to his face as he tugs and fiddles with the bandages holding it in place trying to get it off. The nurse comes into the bay with the mobile medicines station, walks past him and goes over to the man at the end of the bay who is lying in bed and reading the paper. As she is doing this, this man now has the IV tube in his hands and is now pulling at the tubing from the mobile drip stand and is feeling along the length and coiling it up in his hands. He continues to pull at the tubing. [Site B day 4]

This behaviour was typically interpreted by staff as a feature of a dementia diagnosis and interpreted as the person lacking capacity. As in the cases above and below, the key response was replacement and securing and tightening the equipment in place. We observed, few (if any) attempts to consider if there was an underlying reason for this, such as the person not recognising what is attached to them, why it is attached to them, or that it may be uncomfortable or causing pain or distress:

This woman, who is 87 years old, has vascular dementia and has been admitted following a fall and has a minor head injury (stitches on face) and fractured right wrist, is sitting (perched on a large pillow) in the chair at the side of her bed, wearing a pink hospital gown, grey pressure socks and black leather shoes. She calls me over and tells me she is very unhappy: I am not happy, I don't want to be here, my arm is hurting me. She rubs her tiny thin arm, the IV port in her arm is secured in place with a huge amount of white bandage wound tightly around it, presumably to keep it in place, it covers almost all of her lower arm. She rubs her arm and I can see the very delicate and thin skin on her upper arm is covered in dark purple bruises. The trolley is in front of her with her lunch, scampi, peas, and chips, which is untouched. [Site E day 4]

This type of resistance often interfered or interrupted with processes identified by staff as critical to the patients care and could delay discharge or transfer to a more appropriate setting, which could be particularly frustrating for staff. The following example is of a man living with dementia admitted to MAU who needed a blood transfusion before he could be discharged back to the care home where he was living. He repeatedly removed the cannula in his arm and allowed the team to repeatedly reconnect it and re-start the blood transfusion he needs. He becomes increasingly frustrated with being kept on the ward:

I have a conversation with the SHO regards this patient, who has again removed his cannula so he can go and put his uneaten sandwich in the bin. She tells me that you just have to keep retrying and hope he stays distracted long enough to manage it. She says she will not restrain him as she believes this is morally unacceptable, she also will not classify him as refusing the cannula as he always allows them to (re)connect it and consents to his treatment, he just later forgets what it is and pulls it out, she says they are: running out of places to put it. What do you do if a patient needs blood but won't keep the cannula in? Keep trying till I get pissed off and pass it on to some other poor sod to try! [Site C Day 9]

Removal of sheets and clothing

Although the removal of bed sheets by a patient may appear to be a relatively non-problematic form of resistance in the context of the delivery of care, it could have wider impacts on how the person was viewed by staff. The removal of bed sheets often resulted in the patient exposing their genitalia or continence pad to staff or to the wider bay, was highly visible to staff (across all sites).

Unlike shouting or crying, the removal of bedsheets and subsequent exposure would always be immediately corrected, the sheet replaced and the patient covered by either the nurse or HCA present on the ward. The act of removal was typically interpreted by staff as a feature of dementia and staff responses were framed as an issue of patient dignity, or the dignity and embarrassment of other patients and visitors to the ward. However, the response to removal was always the replacement of the sheet and this could trigger further cycles of removal and replacement, leading to escalation of other behaviours associated with resistance.

An example of this is a go year old woman admitted to site C. She has been admitted to the medical assessment unit despite having no immediate medical need apart from her dementia diagnosis. Instead she has been admitted from a care home, where she has been placed as a result of her husband having a stroke and no longer being able to care for her. Her behaviour over the previous evening and morning has involved shouting and refusing all food and care and has required assistance from the specialist Dementia Care Worker. However, she has been calm since a visit from her husband earlier in the day, and has eaten. The care home will not readmit her, so she is now stuck on the unit (in this case an overflow unit due to a high number of admissions to the Emergency Department during a patch of exceptionally hot weather) until an alternative appropriate destination can be found for her. During observations, she remains calm for the first two hours. When she does talk she is very loud and high pitched, but this is normal for her and not a sign of distress. For staff on the bay their attention is elsewhere, because of the other 6 patients on the unit, one is on suicide watch and another is refusing their medication (but does not have a diagnosis of dementia). At 15:10 she begins to remove her sheets:

15:10: This patient has begun to loudly drum her fingers on the tray table and has still has not been brought more milk as requested from the HCA an hour earlier. The unit seems chaotic today. As it is a temporary overflow unit staff do not know where things are, noticeably there are no cupboards or units. She has moved her sheets off her legs, bare knees peeking out over the top of piled sheets.

15:15: The nurse in charge says hello when she walks past her bed, and she looks and smiles back at her. She explains to her that she needs to shuffle up the bed and she asks about her husband, and reminds her that her husband was there this morning and that he is coming back tomorrow. However, she does not believe the nurse who tries to reassure her that he has, but she sounds upset.

15:20: She now asks every passing nurse or HCA if she can go, and it sounds as if she is pleading because her voice is so high.

15:25: I overhear the nurse question, under her breath to herself, why this patient has been left on the unit and she has started asking for somebody to come and see her. The nurse tells her that she needs to do some jobs first and then will come and talk to her. The lady in the next bed reassures her, tells the nurse she will talk to her instead. The patient goes quiet.

15:30: She has once again kicked her sheets off her legs. A social worker comes on to the unit looking for a different patient and the patient calls her over . The social worker replies 'Sorry I'm not staff, I don't work here' and leaves the bay.

15:40: She keeps kicking sheets off her bed, and she now whimpers whenever anyone passes her bed, which is whenever anyone comes through the unit's door. She stops whenever the neighbouring patient reassures her. She is the only elderly patient on the unit and again the nurse is heard sympathizing that this is not the right place for her to be left. [...] this continues and eventually she starts sobbing, getting more and more upset as she talks, the ward staff keep talking about husband and home, which is where she wants to be and the one place that she cannot go. [Site C MAU, Day 13]

Of note is that the unit was uncomfortably hot and stuffy, and a need to be uncovered and cooled down could be viewed as reasonable, and in fact was considered acceptable for other patients (without a classification of dementia) provided they were otherwise clothed. This is an example of an aspect of care where the choice and autonomy granted to patients assesses as having capacity is not available to people who are considered to lack capacity (classified as having dementia) and carries the additional moral judgements of the appropriateness of behaviour and bodily exposure. In the example given above, the actions were linked to the patient's resistance to their admission to the hospital, driven by her desire to return home and to be with her husband. Throughout observations over this period patients perceived by staff as rational agents were allowed to strip down bedding for comfort, whereas for people living with dementia, this was often interpreted as a deviant behaviour, and would be challenged and corrected by staff.

Language & performance: communicating the 'rules' of the ward in response to resistance

In the provision of care at the bedside and in response to perceived resistance, the ways in which ward staff talked to patients living with dementia was highly repetitive. It typically addressed the person by locating them very clearly in relation to the reality of where they were, what had happened to them and what was going on around them. This talk focussed on reorienting and locating the person to the rules of the ward and the accepted behaviour within it, and overcoming perceived resistance to care or to manage and contain the person within the context of the routines of the ward. However, people living with dementia did not learn the rules and fit into the timetables of the ward. Importantly, this talk was rhetorical and did not require or expect the person to respond or assent. The content of this talk did not necessarily reflect what staff actually did, with care and work on the body continuing during these encounters.

The most common approach staff used in response to resistance was to locate the person living with dementia within the institution 'you are in hospital', and the reality of what had happened to them 'you have broken your hip'. Staff also gave very clear instructions to be followed and obeyed, often emphasising the potential imminent danger of a patient's actions and these typically contained a powerful sense of urgency that often displayed their own underlying anxiety and fears. Staff also negotiated and bargained with patients offering to leave them alone and stop disturbing them if they cooperated with the request. Appeals to the necessity and expectations of the institution was commonly referred to, and these appeals emphasised that there was no choice for either the person or the ward team caring for them 'we have to change you'. This repetitive talk was directed at reminding the person living with dementia of their place in the world and the status of ward staff, they must all fit the expectations and timetables of the institution.

These exchanges provide ways of uncovering aspects of the loss of identity and social standing of people living with dementia. It is important to note how much of this talk is directed at trying to remind the patient of their place in the hierarchy of the ward and the rules to be followed. Although staff often gave the appearance of seeking permission and negotiating with the person, the delivery of care typically continued as staff carried out this 'talk', with a tacit assumption of assent, focussed on obtaining the correct response from the person to allow care, which was already being carried out, to continue.

Use of multiple interactional approaches

Importantly, this talk was repetitive and cyclical. Although orienting people to the reality of where they were 'you are in hospital' and what had happened to them 'you have broken your hip' were the most common ways in which staff responded to resistance, they typically employed a cascade of techniques as each in turn failed to obtain the appropriate or required response: the patient's acceptance of their request and to allow the delivery of care to continue.

Here, the team use a number of different techniques with this 95 year old man with a diagnosis of dementia who has been admitted with a fractured hip: they start by asking this person if he wants to get out of bed, they emphasise the necessity of this, and then appeal to the requirements of others (physiotherapy team and his family). Throughout this encounter the team discuss together what they are doing, emphasise his autonomy and his ability to decide 'I won't force him'. However, as they talk they start to work on the patient's body. When they complete this task and he is sitting in his chair, they praise him and reward him with a chocolate.

He is lying in bed and the bed is very low to the ground, the nursing team go over to him and ask him if he wants to get out of bed.

Patient: no leave me here

The team remind him that his daughter will be here soon: you were going to walk with the physio for her, you need to be up in the chair when your daughter arrives, your family will be with you soon and they will like to see you up in the chair, are you sure? I am not going to force you but it would be good to have you sitting up. Lets make a deal, we will give you this morning to rest and get you up later in the afternoon. We just need to check your pad. Patient: I am not well...

HCA: We still need to check the pad for you, we must get you up this afternoon. They discuss together that they won't force him: it's not fair, he can still tell you what he wants, it's not fair. They draw the curtains back and sort out the bedside around him and tidy up the trolley, and give him some chocolate and pass him some water with a straw and he drinks it. The physiotherapist comes over to chat to the team and they discuss him, they have arranged to work with him and get him standing when his daughter arrives at 1pm: Your daughter is coming at 1 o'clock so we can do some walking, will you let the nurses help you up. He is clearly not keen and they remind him: you broke your leg a few weeks ago. I will see you at 1pm. She kneels at the side of the bed and leans over the bed to him. She has a very kind tone. Later they go back over to him and wake him up: Sorry my darling, we are going to get you up in your chair for your lunch. As we discussed my darling, your daughter will be here. She bends low chatting to him in a very friendly and chatty tone and they draw the curtains: We are going to get you up

Patient: Can I stay here? He sounds frail and very pitiful: can't I stay here, don't make me get out of bed, my back aches

Team: We spoke about this, you need to get in the chair, you are doing some exercises, your back probably aches from lying in bed

Patient: I'm going to die

Team: You're not going to die, can you roll that way or lift your bum please, you liked to be in your chair yesterday, can you roll on your side

Patient: It hurts

Team: Well done, we know, we are being as guick as we can

Patient: Sorry I am like this

Team: It's not your fault, well done! There we are well done, just relax, pop your hands on your chest for me. They are using the hoist suspended from the ceiling- I can hear it buzz. Put your hands on your chest and we will do the rest my darling, that's it well done, there we go, see not too bad. I know, we will be as quick as we can, we will get you in the chair as soon as we can. They position him in the chair: there we are, don't you feel better sitting in the chair? For all that hard work would you like a chocolate? I thought that would make you feel better, she gives him a chocolate from the bowl on his trolley. [Site C day 6]

It is important to consider the intensity of this work. Given that all the work of the wards and routines, which are not bundled together but are separated out into individual timetabled tasks, each provided by different members of staff. This meant that these exchanges were a recurrent

feature of the timetables of the ward and repeated again and again at each bedside throughout a shift.

Orienting to the reality of their situation

A key approach staff used to communicate with people living with dementia when faced with resistance was typically by locating them very clearly in relation to the reality of where they were, what had happened to them and what was going on around them. This was the most commonly used approach, consistent across institution, as the most frequently and universally utilised approach when faced with resistance. Only within two MAU units (sites D and E) did staff consistently not use this approach with patients.

Typically, these rationalizing statements placed an emphasis on the reality of where the person was 'you are in hospital', that they cannot go home, provided details of their condition 'you have broken your hip' or 'you have an infection', and directly contradicted patient's statements or their perceived reality, such as 'there are no policemen'. During these encounters, ward staff appeared to be actively trying to support and help orient the patient to the reality of where they were and what was happening. However, this approach always appeared to increase the person's anxieties and concerns and triggered further resistance. In this example, the team either only briefly acknowledged or did not respond to the anxieties this 87 year old woman with vascular dementia who has been admitted following a fall, a minor head injury (stitches on face) and fractured right wrist expressed throughout their encounter (wanting to go home, the cost of hospitalisation and where her family is). Instead, they focussed on repeatedly reminding her of where she was, 'you are in hospital':

HCA: do you want to go to bed? Patient: NO, I want to go home

HCA: You are in hospital, you are in hospital

Patient: I can't afford it

HCA: It's free

She rubs her arm with the bandaged IV port and runs her hand along the long tubing leading to the mobile stand and the HCA states that this is: for your medicine

Patient: I don't like being here I don't know it

HCA: I know, you are in hospital. She puts the orange juice pot from lunch into her sip cup and puts it in front of her

Patient: Where is my son

HCA: He will be here soon. The physio team arrive and the young female physio stands next to her while the other gets a walking frame from the bed opposite and brings it across.

Patient: I am NOT staying here

Physiotherapist: She crouches down beside her and looks up at her: at the moment you are in hospital, you are in hospital. You have hurt your hip. She is sitting slightly propped up and lying on one side and she is hanging onto the side rails and she is looking very agitated again.

Patient: where is my son?

Physiotherapist: I am sure he is coming later, would you like your cardigan? She is just sitting in her hospital pink gown and now they have mentioned it, she does look a little cold. They get it from the cabinet and help her to put it on and she instantly looks more

comfortable. They ask her: What country are we in?

Patient: I don't know, I can't afford to pay for it anyway. The physio team decide to give up and to leave her and they move on. [Site E day 4]

This approach also extended to family, who also typically repeatedly reminded the person of the reality of their situation.

Clear instructions to be obeyed

Ward staff also typically gave people very clear instructions to be obeyed, often emphasising the potential risks and imminent danger of their actions if the person living with dementia did not comply. These typically contained a powerful sense of urgency that often displayed their own underlying anxiety and fear. These exchanges provide ways of seeing the loss of identity and social standing of people with dementia within the ward. As in this case, staff often raised their voices as they gave very clear and often very pared down and simplified instructions to the person. Here, this HCA is providing one to one care for a 74 year old man who has a diagnosis of dementia and has been admitted with a chronic subdural hematoma:

The HCA is talking in a very loud voice to him: EAT YOUR BREAKFAST SITTING DOWN, NOT STANDING.... EAT YOUR BREAKFAST SITTING DOWN, NOT STANDING. He stands over the patient as he is sitting on the bed. He is in pyjamas with a beige jumper over the top and the pyjama bottoms are far too long and are puddling around his feet on the floor. He is staring unfocussed in front of him with a very blank expression on his face and although the HCA is shouting at him, he does not appear to respond to what is going around him or to the team. The mobile trolley is in front of him with his breakfast and he has buttered his own toast, he eats a bit of this and also quite a bit of his rice crispies. The HCA sits in a chair at the door and occasionally as the patient starts to stand as he eats his toast, the HCA goes over to him and tells him: EAT YOUR BREAKFAST SITTING DOWN, NOT STANDING. After breakfast they walk together down the corridor to the day room and back again. When they get back to the single room, the HCA says: SIT IN THE CHAIR, SIT DOWN

He does not respond and stands still in the middle of the room and the HCA repeatedly tells him: YOU ARE IN HOSPITAL....YOU ARE IN HOSPITAL....YOU ARE IN HOSPITAL....THERE IS NO GIN AND TONIC, SORRY MATE.[Site B day 3]

This demonstrates how repetitive behaviour is viewed as particularly problematic and becomes the focus of care. In the examples above, repeated attempts to stand becomes an urgent focus of control. These rationalisations also typically included repeated warnings of danger and the likely consequences of their behaviour, if they continued and do not modify their behaviour, in this case, the common fear amongst ward staff of people living with dementia being at high risk of falling.

Appeal to the necessity of the institution

Appeals to the necessity and expectations of the institution was commonly used to persuade people living with dementia to accept care, and this typically emphasised that there was no choice for either the staff members or the person, for example, 'we have to change you'. It is important to note how much of this talk took place as staff were already delivering this care and working on the patient's body. This is directed at trying to remind the patient of their own status and the status of ward staff caring for them, they must all fit in to meet the expectations and rules of the institution.

In this case, the HCA initially asks the nurse in the bay to help her to change the wet soiled sheets. She emphasises 'we need to move you' and 'we can't leave you' while the patient remains lying in bed (a 94 year old man admitted with a fractured hip and pneumonia) and is shouting '999', which may indicate that he sees this as an assault or attack. The shouting alerts the wider team and another HCA joins them behind the curtain to provide additional help as they struggle behind the curtain.

The nurse and HCA are back at his bedside 'sorry' they use the shortened version of his name and keep forgetting his first name.

Patient: GET OUT GET OUT

The nurse talks to the HCA: he's soaking, we are going to have to change his sheets, and then to the patient: 'you are wet'. The nurse asks for more help from staff in the ward to help them to roll him and then goes back to the patient: we need to move you, its not good to lie in a wet bed' and in response he shouts: 999 YOU CAN LEAVE ME ANYWHERE YOU LIKE. Nurse: come on now we can't leave you in here, we are just going to change the sheet, you need a new sheet. He is clearly struggling with them behind the curtain and is clearly not happy and his cries and groans can be heard from outside of the curtain as they change the sheets. They draw back the curtains and the patient looks very small in the bed with his head resting on large white pillows. He is now wearing a blue hospital gown and the sheet and a thin blue blanket are tightly tucked in around him with the side bars of the bed raised. He doesn't move. [Site A day 4]

Despite the patient's verbal and physical resistance to the immediate task in hand, once this has been completed the patient is almost instantly settled. This can also be a feature of resistance to care, it can sometimes appear to be limited to the immediate and specific task in hand.

Here, the ward team approach the person emphasising that they need to 'clean' her and need her to keep covered to fulfil the requirements and rules of the ward. As they deliver this personal care behind the screen the team emphasise that they 'have to' clean her. The patient who is a 96 year old woman living with dementia who has been admitted with a fractured hip cries out 'they are hurting me' asking for help 'OH PLEASE HELP ME': The bay team have pulled the screens around her bedside and sound as though they are changing her clothes and the bed sheets. As they do this she cries out: OW, it hurts. The team reassure her as they work: We have to clean you, you are alright you are not falling, we need to clean you, as she continues to cry out: Please, please, help I am getting hurt, its hurting, HELP ME, OH PLEASE HELP ME, HELP ME, HELP ME oooowOW and this ends with a piercing scream. The team reassure her: You are not falling, and have drawn the curtains back and take large piles of linen away in a bag. The patient is a tiny figure in the large bed, covered in fresh sheets and a blanket. Immediately she pulls off the blanket and sheet from her legs- she exposes her tiny legs that are both covered in thick bandages up to the knees. The tea trolley arrives and the young man with it says: cup of tea darling? In response she pleads for help: Please, please, help me [Site E day 8]

Care that is seen by the ward staff as essential, typically continued despite patient's resistance, expressed either physically (pushing staff away) or verbally (crying out). Rather than walking away and leaving the patient, resistance to everyday routine care often resulted in staff continuing with the task while also emphasising the necessity and expectations of the institution by repeatedly emphasising to the patient that they must accept care. Typically, other team members within the ward would be called on to help and provide support or who react to the disturbance and noise by leaving their work (there is rarely a moment when a member of staff is ever inactive and obviously between work or not actively working at the bedside or completing paperwork) to support and help complete the delivery of care.

Negotiation and bargaining

Staff also negotiated and bargained with patients. This was often an approach used when all previous techniques and approaches had been repeated, failed and staff appeared to be becoming tired. With the nurse leading the medication round often negotiating with patients that if they

cooperated with the request, then they would leave them alone and stop disturbing them. Here, as the nurse dispenses the medication, he reminds the person, who is a 98 year old woman with a diagnosis of Alzheimer's and admitted following a fall, of where she is, emphasising the institutional requirements and negotiating with her to complete care:

The nurse is at the bedside for the medication round and asks her: what is your birthday? when is your birthday? Do you remember your birthday? she wakes up and listens and points to the wall. Do you want your paracetamol? So just a couple of tablets? one at a time?

He stands at the side of the bed, passes her the tablets one at a time followed by a glass of water. She tells him she wants to go home. Nurse:

Yes how far is home? You are in hospital now, this is one the doctors asked me to give you. He puts a tablet in her mouth and holds a cup of water while she sips on a straw, but she takes it out. He repeats: You need to take that one darling you've got to take it my darling, pop it into your mouth, I will wipe your hands for you, take a drink and try to wash it down for me. I tell you what, once you have taken these tablets you can shut your yes and I can leave you alone for a while [Site C day 2]

Family and carers responses to resistance

Family members typically struggled with resistance and found this both embarrassing and stigmatising within the ward. This was particularly the case if the person's diagnosis of dementia was recent (a diagnosis was often received during an admission) or if their condition had deteriorated or changed significantly during admission. In response, families also typically attempted to rationalise with the person, contain them at the bedside, or limit their behaviour in some way. They were also very likely to apologise to staff. This person has a diagnosis of dementia and has been aggressively refusing medication and personal care all day. It has taken up to five members of nursing staff to change her clothes and bed sheets, and they continued as she carried on shouting, swearing, spitting, biting and scratching. Of note is how upsetting this behaviour is viewed by her adult children, but also how embarrassed they are by it.

15:45: 2 visitors are with her. Daughter and son in law. I speak to them and take verbal consent to observe them and the RN comes over and explains her history since admission. Concerns have been raised about her taking medication. Because of her aggression they have resorted to crushing up her medication and feeding them to her in her porridge. The visitors are shocked to hear that she has been biting the staff but say that shouting is not out of the ordinary. They say her standard voice is the loud one, but she still often demonstrates a sense of humour, and puts on a bad northern accent. They are not surprised she is refusing medication and are impressed staff have had any success. They warn the nurse that she can be very strong. A doctor then comes over, the RN leaves, and the doctor begins to discuss her condition with them. They stand at the foot of her bed, discussing her while she sleeps. While the nurse discussed social and behavioural aspects of care the doctor is much more clinical. Later I speak to her daughter. She has been crying and is embarrassed to hear about the biting but says at the same time it is a relief because yesterday she was barely moving - at least the agitation shows that she is awake [Site D, Day 9]

Such response were not uncommon, and across all 5 sites families of patients who were shouting, 'wandering' or refusing food, to instruct their relatives to fit in with the routines of the ward and 'behave themselves'. Their response in part is similar to the parents of a naughty child, in that while what would be construed as bad behaviour is understandable to them, it is also embarrassing (courtesy stigma). This is associated with the wider moral assessment often underlying staff approaches to resistance.

The moral classifications of resistance

We found that people living with dementia who resisted care, particularly when this was regularly disruptive to the ward routines and timetables were vulnerable to moral assessment. Their resistance was often interpreted by staff as another feature of who the person was and this could become a key part of their identity in the context of the ward. In addition, some patients were also identified by the ward team as more likely to resist care based on their assessment of their personality and background.

Judgements of individuals: resistance as identity

While resistance was seen by ward staff as a feature of dementia, staff also judged longer-term, chronic and disruptive resistance as another feature of who the person was and this could become their identity in the context of the ward. In a small number of cases this could take the form of affection from staff: 'she's alright, she just likes a bit of a strop' [Site B day 15] 'she does make me laugh though' [Site D day 8], but more often it was viewed negatively: 'he is a puncher' [Site C day 2] 'she is a hitter' [Site E day 5] or 'a climber' [Site A day 18]. Such people were typically approached with caution, but for staff, there was always an underlying expectation that a patient resisting care could respond physically. Some, usually male (although not always), patients would be approached by staff in pairs, or a male member of staff would be asked to approach them, regardless of their present mood or behaviour.

Some people were also identified by the team as more likely to resist care based on their assessment of their personality and background. Here, the team discuss a 94 year old man with a diagnosis of dementia admitted with a hip fracture and pneumonia, who had resisted care the previous night and they rationalize that this person's character and heritage means that he is likely to resist further care. In response, they decide to approach him with caution. Two members of staff approach his bedside and focus on asking his permission, standing at either side, holding his hands and as they gently talk to him, checking whether he prefers that they use the full or shortened version of his first name and check his 'this is me' document (This is a form to support 'person-centred care' and includes space for families to provide details on the person's cultural and family background; key events, people and places from their lives; preferences, routines and their personality). He resists their request by shouting 'NO'.

All is quiet in the bay. The nurse talks to him: are you cold? You sound chesty? The bay team and the anaesthetist discuss him: when he came to us from A&E, he was fine yesterday. Chesty, it started overnight. He was fine yesterday, I don't' think we should do him, he's strong willed, Irish people. The HCA and the nurse are on either side of him and hold a hand each. Can I take your temperature? Patient: NO, MIXED UP. He pulls his IV out of his arm and they try to take his temperature. They ask him his name and whether he prefers his full name or a shortened version and they refer to his 'this is me' document. [Site A day 4]

Resistance as deliberate and wilful

Resistance could also be interpreted by staff as an act of wilfulness to express their responses to their hospitalisation. This in turn, overshadowed the person and any opportunities to identify any potential underlying patient need. For example, this 94 year old man admitted with a UTI and has 'acute confusion' is in a single room and regularly cried out for help and had fallen a number of times during his admission. The repetitive nature of this behaviour meant that he is characterized by ward staff as deliberately acting in this way to disrupt the ward and their work. The ward sister rushes to the room and expresses her frustration and shouts 'WHAT ARE YOU DOING' at him.

The ward sister tells us he is in pain, but he knows what he is doing, he says 'I will shout as much as I want'! I asked the doctor for some diazepam to calm him down. Later there is a crash and cry near the entrance to the ward. The ward sister immediately pops her head out of one of the bays and asks me- is that in this ward? She runs into his room: WHAT ARE YOU DOING! She shouts for everyone and they all go running down the corridor and all are with him. [Site A day 10]

In some cases, although resistance could be interpreted by staff as a deliberate choice, they also attached wider emotional meaning to this. During the shift handover, the team discuss a patient who is resisting all care 'he is declining everything'. They suggest that although he is 'confused', the underlying reason for this behaviour is that he has 'given up' and does not want to live.

I am immediately struck that they are doing a handover at the bedside and the details of patients on the handover sheet, this is incredibly detailed for each patient. I follow the team and they head to the bay at the far end of the ward, there are 5 nurses including the handover person, although I am a bit overwhelmed because so much is going on. One of the team is pushing the mobile computer station and is updating the notes as we go through them. The team help me to find the right page- we are working back from the end to the beginning of the handover sheet and we go from bed to bed. We stop at the bedside of an 83 year old man who has a diagnosis of Alzheimer's admitted following a fall and 'general decline'. The team discuss him: He was very confused overnight, declined everything, I think he has given up, very confused, he was fighting with us. [Site B day 1]

This is associated with staff beliefs about the potential future and quality of life for people who have dementia.

The emotional and somatic impacts of the organisation and timetables of the ward Our observations identified that resistance appeared to be a response and reaction to the impacts of an admission on individuals. These were both emotional and somatic and included the difficulties communicating needs, high levels of anxiety, and the unfamiliar environment of the ward, which could lead to disorientation. A key impact of this was the person being viewed within the ward as having increased dependency, which in turn resulted in them loosing skills and independence.

Resistance was typically triggered by the unfamiliarity with the ward, or the fixed routines, timetables and the organisation and delivery of care within an acute ward itself. Loud noises, bright lights, and unfamiliar people were distressing for many of the people living with dementia observed. Because care in these settings is delivered by shift and rota there was little continuity of care, exacerbating the unfamiliarity of the environment and the wider distress of other patients within it.

As such, resistance can be framed as a rational response to the organisation and delivery of care, with routinized, timetabled and compartmentalised care producing fear and anxiety within a patient, but never acknowledged in the assignment of staff at the handover of each shift. Our analysis, drawn from observations that allowed the researchers to spend extended time within and across shifts observing specific ward bays and the individual staff and patients within them. This observation almost always revealed the potential underlying reasons or triggers for patient's resistance to aspects of care or their admission. These were typically rational to that patient's present ontology and perceptions.

Difficulties communicating care needs

Trying to stand, standing, walking, agitation, or being unsettled in bed or at the bedside was typically interpreted by ward staff and responded to as resistance to care. The most typical initial response from ward staff was to return people to the bedside, encourage them to remain in the bed or the chair and then to repeatedly try to contain them at the bedside. There were common patterns across all sites of an agitated patient standing and being repeatedly returned to the bed or chair before staff recognized an underlying care need or the patient was eventually able to communicate verbally that they needed care, such as continence care and wanting to go to the bathroom.

Trying to stand, standing, walking, agitation, or being unsettled, appeared to be a key and common form of communication and a sign that the person had an underlying care need. For this group, who often had difficulty articulating and communicating their needs verbally to someone, this usually involved often extended periods of staff returning people to their bed or chair before ward staff recognised there was an underlying care need. However, this recognition did not always occur. This was often associated with people trying to go to the bathroom or requiring help with continence care. Here, the immediate staff response was to return this 93 year old man who has a diagnosis of dementia and has been admitted following a high number of falls to their chair and it took time and repetition for their underlying care need to be recognised:

The HCA remakes the sheets of the neighbouring bed and while she does this he starts to stand up, holding onto the slides of his chair She sits him down. The HCA is focused on his neighbour and asking him what he had for breakfast and he asks for more tinned fruit, there is a tin open on his trolley. As she does this, the 93 year old man stands up and the HCA goes and gets him to sit down and he says he wants the bathroom. [Site A day 6]

However, even if a person living with dementia did display the signs and cues in their body that something was wrong and that they were seeking attention and help, this could be a low priority for ward staff. In this example, this 96 year old woman with a diagnosis of dementia admitted with a fractured hip, became increasingly anxious and distressed, which can be seen from her fiddling with and pulling at the band of her skirt, and talking in an increasingly animated way, and trying to get attention from passing ward staff. However, even though this seems to be causing her great anxiety, this appeared to be a low priority for the ward team particularly as in this case, when there was no immediate risk to the person (she safely contained in the bedside chair) or perceived urgency if the person is wearing a full continence pad (or a catheter) as is the case for this woman.

She now looks very distressed, she is talking to herself and looking around the room wide eyed and fiddling with her skirt. Everyone is busy, the nurse is focussing on the drug round and no one else is here, so I go over to see her and she reaches out and holds my hand. Her speech is very hard to understand, but she tells me that her leg hurts and she touches her knee. I tell the nurse in the bay and she says they will put her back to bed soon and check on her. I go back to the bedside or let her know and she keeps me at her side and says clearly: can you take me to the toilet? Can you take me? Can you take me? I say I will tell the nurse and I do that again, and she responds that she will check her pad when they put her to bed soon. Some time passes and she is now fiddling with her skirt even more, no one has come after I alerted the nurse so I go over to the male HCA and tell him and he responds: I will get the hoist to put her back into bed and I will check hr pad then but she fights! He smiles. Later the bay team use the hoist to put her back into the bed and she becomes very upset. [Site E day 4]

This woman had also been described by ward staff as difficult to care for and as someone who 'fights' staff providing care. This exacerbates the delay she experienced because they decide to

bundle her care (personal care and toileting) together. However, by the time the team were ready and have all the equipment (the hoist must be found and brought from the other end of the ward) at the bedside, the patient is extremely distressed and although the team asked her why she is upset, they did not connect this to the delay she has experienced and her underlying anxiety of losing her continence.

Unacknowledged anxiety and the impact of an unfamiliar environment and routine People living with dementia appeared to have high levels of anxiety throughout their admission, with high levels of underlying anxiety observable in every person living with dementia within the study at some stage during their admission. This included anxieties about where they were and what was happening to them. A key anxiety was not knowing when they were going home, whether they would be able to leave and go home, and if they still had a home. This 81 year old woman with a diagnosis of dementia, who had been admitted with a fractured hip, but was now medically fit to leave, woke up during the day and in a very anxious and trembling voice asked me and the HCA a large number of questions that displayed the uncertainty she was feeling about what would happen to her:

She wakes up and the HCA goes over to her and checks on her, she is very gentle and props her up in the bed and gives her a sip cup of juice and she takes a drink. She sees me and calls me over and take her hand and she looks and sounds quite anxious and asks in a tiny trembling voice: How long have I been here? When can I go home? I will get well wont I? Do I have a home to go to? Have I been here long? When can I go home? I will get well wont I? Do I have a home to go to? Have I been here long? When can I go home? Why am I so tired? I will get well wont I? Do I have a home to go to? Have I been here long? When can I go home? I will get well wont I? Do I have a home to go to? Why am I so tired? I hold her hand and reassure her and tell her she is safe and I give her the sip cup to drink and she takes a sip and I encourage her to drink a bit more. She has a nasty chesty cough that she didn't have the day before, which worries me. She is lying slightly propped up in bed covered in a blue blanket tucked around her feet. I suggest she closes her eyes and that I will stay here until she falls asleep. She falls asleep straight away. [Site E day 13]

In addition, a person's anxiety about where they were and what was happening to them could also quickly heighten, if the unfamiliarity of their experiences increased in any way. Here this same 81 year old woman above becomes increasingly distressed during a shift. She has no clean clothes left so the ward staff dressed her in a hospital gown. However, for this person, not wearing her own clothes, particularly, for her, not wearing her bra and trousers, were essential garments, and being without them appeared to contribute to her becoming increasingly anxious and upset. It increases her confusion about what is happening to her, where she is, and makes her feel that she has 'lost' her clothes. It also confirms to her that she is somewhere unfamiliar and emphasises that she cannot leave.

I arrive at the ward and go over to say hello and she says: I still love you. The one to one HCA is with her tells me that she is very teary and upset today, that she has just had a shower and she doesn't have any fresh clothes and so she is now in a hospital gown. Patient: I want my trousers, where is my bra I've got no bra on. She is very teary and upset, she is sitting in a pink hospital gown and her lilac cardigan on over it and it is clear she doesn't feel right without her clothes. The HCA asks her: Your bra is dirty do you want to wear that? Patient: No I want a clean one. Where are my trousers? I want them, I've lost them. The HCA explains that her clothes are dirty and this discussion continues for some time. The ward cleaner arrives to sweep around her and as he does this is says hello to her and she tearfully explains that she has lost her clothes and he listens and is sympathetic. She repeats: I am all confused, I have lost my clothes, I am all confused, How am I going to go to the shops with

no clothes on! Will I get out of here? She is very teary and upset. I sit next to her and hold her hand and the therapy dog arrives to see her. [Site E day 5]

Disorientation-place and time

People living with dementia could become quickly disoriented; sleeping patterns could become altered by the acute condition, their treatment, the ward environment and by the fixed routines of the ward. Sleeping during the day (although staff talked about the importance of discouraging daytime sleeping, in practice leaving people living with dementia to sleep during the day was normal practice across all sites), and the phenomenon of 'sundowning', was a pattern of behaviour widely reported and recognized by ward staff where people became increasingly agitated and active at the start of the night shift and did not sleep during the night. In addition, during night shifts, the routine work of ward staff, which involved delivering care to the patient at the bedside continued. During the night shifts, many people living with dementia believed they were in their bed at home and so ward routines such as personal care, medication, and turning people in their beds (to prevent pressure sores), which was typically carried out in the dark to people living with dementia who were asleep or semi-conscious could further disorient them and increase their anxiety and fear. This meant that the night shift could be a very frightening time for people living with dementia and even for people without cognitive impairment, it could be disorienting. This 85 year old man who was admitted with a fractured hip, does not have dementia, and is blind and so the experience of the ward team at the bedside was very frightening and disorienting:

The team are at the nurses station discussing what happened during the night shift. They chuckle as they tell me that in the night a male nurse and a male HCA were caring for him and he through they were attacking him. He said he was going to call the police, but they didn't think anything of it, but he used his mobile phone and called the police and told them that someone was attacking him in his own home. They went round to his house and eventually they called the ward. The laugh and tell me it happens sometimes. Later that day he calls me over to his bedside and tells me: I am so upset with myself after all the care they have been giving to me, to sort of throw it back in their face. It was so real to me, it was dark and I really thought I was in my own home and I could just see shadowy figures and I called the police, I am so sorry I am ashamed. He starts to cry and big tears roll down his face, I give him a tissue and he dabs his eyes and screws it up in his hands. I hold his hand and reassure him and he asks me: can you apologise to the nurses for me? I pass this on to the nurses and return to his bedside and we discuss his travels around the world, we talk about Rio and his experience of sailing around South America and Cape Horn. [Site C day 8]

Although this was not seen by ward staff as unusual, the additional feature of it leading to the involvement of the police meant that it became an amusing story that circulated quickly around the ward. However, this could have longer term consequences for the person and their sense of self.

Ward strategies of care in response to resistance

Overall, the key response to resistance and refusal of care by people living with dementia was one of containment and restraint. Although specific techniques had some variance between wards, the overall strategy was always to keep the person living with dementia within their bed or sitting at the bedside. Across all sites, staff expressed high levels of concern and anxiety about people living with dementia attempting to or leaving the bed or bedside, and this increased exponentially if they were standing, walking in the bay, the wider ward and corridor or close to the ward entrance. Multiple techniques of containment and restraint were observed being used by ward teams. Importantly, these approaches to patient care and their containment at the bedside was both a response to resistance, but were also frequently the trigger of resistance or cause of patient anxiety.

In addition, a key response to ongoing identified regular patterns of resistance by individuals living with dementia, within wards was 'specialing' and to assign one-to-one agency HCA staff to care for them. This was used particularly when individuals were disrupting the ward routines and described as a way to support both people living with dementia and ward staff.

Cultures of containment and restraint

The key ward staff response to resistance and refusal of care by people living with dementia was one of containment and restraint. Although specific techniques had some variance between wards, the overall strategy was always to keep the person living with dementia within their bed or sitting at the bedside. Across all sites, staff expressed high levels of concern and anxiety about people attempting to or leaving the bed or bedside, and this increased exponentially if they were walking in the bay, the wider ward and corridor or close to the ward entrance.

Multiple techniques of containment and restraint were observed being used by ward teams. This included raising the side rails of the bed, or tucking bed sheets in tightly around the patient were both common and covert means to contain a patient within the bed. For those patients sitting at the bedside, the close placement of the mobile tray table, unreachable walking frames and technologies such as chair alarms were used to contain people and keep them sitting in their bedside chair. Clinical technologies, including continence technologies (particularly full continence pads, and using bed pans, and commodes at the bedside), medication and sedation, and tightly secured medical equipment also limited or restrained movement from the bedside. Importantly, these approaches to patient care and their containment at the bedside was both a response to resistance, but were also frequently the trigger of resistance or cause of patient anxiety.

Walking away from the bedside or within the ward was overwhelmingly categorised by nurses and HCAs and medical teams as a deviant action. While the question 'where are you going?' would frequently be asked of patients, it would only be done so in a rhetorical sense, a cheery admonishment used as the person living with dementia was led back to their bed or bedside. It was rare for it to be considered that the patient with dementia could be purposeful, with their actions instead always categorised by ward staff as 'wandering'.

With the exception of two closed wards (MAU) where some movement was permitted to patients classified as a low-risk of falling, walking in the ward was always viewed by ward staff as a problematic activity. One closed acute ward also carried out these restrictions on their patients living with dementia, which meant they were not permitted to leave the room or to walk in the corridor. For the person living with dementia, in their temporary role as patient, to be labelled as a 'wanderer' was to be classified as a visible part of the ward, requiring supervision, restraint and control. Such categorisations were made without the assessment or consideration for the patient's mobility or independence of movement prior to their admission. Walking was considered a purposeless activity, with a 'wanderer' seen as drifting on the ward, an activity with little reward but high risk (falls, leaving the ward, approaching other patients). What was not considered is that for the person living with dementia, the activity had purpose. This may have been walking to the toilet, looking for someone or something, or simply to stretch their legs for stimulation. Often, however, they expressed a strong (and arguably rational) desire to leave the ward and to return home. As such, stopping a patient from leaving or 'escaping' the ward was an everyday and common ward activity.

These approaches to containment, through the use of the previously described covert techniques of restraint, coupled with the verbal techniques of orientation and rule repetition often triggered resistance and unhappiness in patients. This led to patients experiencing high levels of anxiety

about what was happening to them and where they were. People living with dementia reported that they felt 'like a prisoner' (Site C), 'feel terrible' (Site B) 'want to go home' (Site E) and were 'unhappy' (Site E) during their stay. Some patients were observed shouting for the police (Site A, B and D) and even surreptitiously using the phone at the nursing station to call their family for help (Site B), because they believed they had been kidnapped or were being held against their will. More often though, the distress of feeling imprisoned by a culture of containment was manifested in the patient's body language. Anxiety was shown through defensive poses such as folded arms, pulled up bedsheets, tightly holding onto bed rails, or nervously staring at doors and people passing by. Here, this 85 year old man with a diagnosis of Alzheimer's who had been admitted with a fractured hip, tells me that he feels 'like a prisoner' and he is very anxious, he cannot make sense of the rules of the ward.

He is wearing hospital green pyjamas and is sitting in the chair at the bedside. When he wakes up and I go over and say hello and he tells me: I am very frustrated and angry, it's like being in a prison here, one minute they say keep drinking and then next they won't let you and no one tells you anything in here, this is like being a prisoner! [Site C day 9]

These techniques of containment and restraint had detrimental consequences to people living with dementia during their admission in terms of both their emotional wellbeing and their physical rehabilitation. The role of physical rehabilitation was often the point where the culture of restraint and containment could become visible to the observer. The therapy team, the physiotherapists and occupational therapists (with crucial decision-making power around discharge routes), can be observed encouraging patients to get up and move around, only for the ward team to immediately approach patients in order to prevent and admonish them for doing so. This 73 year old woman living the dementia and admitted following a fall had refused to carry out her rehabilitation session with the physio team earlier that day and during this shift, the HCA discusses with her and her husband the importance of rehabilitation and attempting the prescribed exercises in order to be fit for discharge. At the same time as these discussions and throughout this shift, she keeps attempting to get out of the bed by putting her legs over the raised side rails of the bed. However, this does not prompt the HCA to help her out of bed or support her walking. Instead each time she does this, the HCAs lifts her legs back over the rails and within the bed. Even when she becomes increasingly restless and agitated, the HCA instructs her to 'stay in bed' and 'try to sleep' and eventually reminds her of the rules of the ward: 'I don't want you to get out, I'll get the sack if you do that', emphasising her interpretation of her role as one of containment and restraint of the patients within the bay.

It is very quiet in the ward, there are a few buzzers going and there are a few members of the team at the nursing station- a mix of ward staff and the medical team. This woman is sitting propped up on the bed, but she looks uncomfortable and she groans. The HCA goes over and puts the side rails up on her bed. She tries to get out and has a leg over the side rails, but the HCA comes over and puts her leg back in: Don't go, you need to be here, lunch is coming soon and then we will get you in a chair, try to sleep. She adjusts the height of the bed and puts the blanket over her and she is now propped up and holds the blanket to her with both hands on it. She is very tiny and wearing a pink jumper, black leggings and red hospital socks which are huge on her, her husband tells me she has size 3 feet. It is warm autumn day and the sun is streaming into the room. All is quiet there is the sound of buzzers outside in the corridor and there is a low hum in the room from the bed pumps. She continues to try to put a leg out of the bed and she tells us: I am not good at just lying here, but I must! She closes her eyes and lies back on the pillows. She again tries to get out of bed and has a leg out over the high sides. The HCA spots this: No stay in bed, and she discusses with her husband that she didn't do the stair exercises with the physio that morning and they encourage her, if she does the exercises that she will be able to go home quicker: you

need to do the exercises so that you can get home. The patient responds: I have no intention of getting anywhere near those stairs! She puts a leg out of the bed again and the HCA repeats: put your leg back in I don't want you to get out, I'll get the sack if you do that. [site 4 day 11]

Across all sites, staff expressed high levels of anxiety about people leaving the bedside, and this strategy of restriction usually involved repeated efforts and staff utilised a range of strategies to limit the person to the bed or the bedside, shadowing them as they walked, barring their way and repeated attempts to encourage or instruct the person to return to the bedside. Any apparent resistance to care typically increased the restriction the person experienced. However, this strategy typically triggered further resistance and increased the person's feelings of anger and anxiety about their situation. It also meant that the person's underlying need may remain unexplored, unidentified and unmet or significantly delayed.

Here, this 81 year old woman living with dementia and who was admitted with a fractured hip, but is now medically fit to leave and is waiting for her 'package of care' is walking away from her bedside. The HCA gives her a walking frame to use and immediately leads her back to the bedside. This restriction makes her very angry. In response, the HCA reminds her that she cannot leave her and shadows her a step behind with her arms crossed, however when she does not comply, continues to walk out of the bay and tries to open the (locked) doors at the end of the ward, the HCA takes the frame from her and turns it around to face into the ward. During this encounter, the HCA starts with emphasising the reality of the ward 'I cant leave you alone', repeatedly reminding her of the rules of the ward 'You need to go back to bed' and then resorts to instructions to be obeyed 'turn around'. However, by the end of this encounter the HCA is very frustrated and this woman is very tearful and upset and is shouting in frustration to be left alone. This also created a lot of attention from other staff in the ward who crowd around her, which further upsets her and adds to her frustration.

She has just woken up and starts to walk away from the bedside. The HCA gives her a walking frame to use from another bedside and stays with her and leads her back to the bedside. Patient: LEAVE ME ALONE, she is very angry and wants to leave. The HCA responds: I cant leave you alone, hold onto it (walking frame) with both hands. As she uses the walking frame to leave the bay, the HCA is staying with her as she goes, shadowing her a step behind, with her arms crossed. The patient turns to her sharply and shouts: LEAVE ME ALONE, she walks up and down the corridor using the walking frame and has arrived at the doors at the end of the bay and she looks through the glass panels and tries to open them (they are locked). The HCA stops her and asks her to turn the frame around and to go back. She then forcefully takes the frame from her and turns it around to face into the ward. This really upsets her: LEAVE ME ALONE...I WISH YOU WOULD LEAVE ME ALONE. I wish she (HCA) would stop following me around! I am not going anywhere! She is very angry and frustrated.

This continues and a member of the physio team returns and says is a very bright and friendly voice: hello! come with me! She puts an arm around her shoulders and steers her down the corridor in the other direction: how about a nice up of tea? The patient now looks very upset now and tearful. [Site E day 2]

This restriction to the bed or at the bedside of people living with dementia extended to older patients, even when ward staff were aware that they needed to be mobilized and required rehabilitation.

Outsourcing care and ward cultures of containment

A key response to an identified pattern of resistance by people living with dementia within wards was to assign one-to-one agency HCA staff to care for them. This was a policy within almost all hospitals and wards as a way to support both people living with dementia and ward staff. In practice, this meant that once a person living with dementia was identified as resisting care within the ward, they could be classified as 'specialed,' a DoLs could be obtained and the ward could legitimately request additional support and assign an agency HCAs to provide one-to-one care to manage that person during their admission.

However, as well as having an impact on the person living with dementia, it could have powerful impacts on the wider ward culture. Ward staff appeared to be less concerned about de-escalating and supporting a person who was resisting care. This request could also function as a means to demonstrate to the wider hospital system that this was a ward and a team that was experiencing specific strains and required additional support staff.

The role of the one-to-one carer was significant in that it was almost always interpreted as a role that required them to restrict the person living with dementia to the bedside, to limit their impact on the wider work of the ward and to enable the ward team to continue their work without interruption. They operationalised their work as primarily involving containment of the person at the bedside and to ensure that they fit within the rules of the ward. In addition, ward sisters reported that although they could request one-to-one HCA care for people living with dementia who were resisting care, they did not believe that they could direct them on how to provide care for the person and that it was not in their power (sites A, B, D). Throughout this shift we can see how both this HCA calls on the requirements and rules of the ward that limit both their work (safety and risk) and the person by locating them within the chair. When this does not happen, both become increasingly anxious and agitated. We can also see here in the language used when talking to this 83 year old man admitted with 'increased confusion', the loud voice shouting, the exaggerated slow pronouncement of words even though they are next to each other in a very small room, indicates how dementia as a condition is believed to affect the person. It is predominantly conceptualised as meaning that repetitive loud and simplified language will eventually enable the person to understand, comply with and to follow the rules of the ward.

He is standing at the door to his single room and using his walking frame, shouting and heading out of the room. The physio and the OT working in the bay go over to him: LETS HAVE A SIT IN THE CHAIR, she speaks in an exaggerated slow and loud voice. He tells them: I want to go home. Physio: YOU CANT LEAVE YOUR ROOM, YOUR BEDROOM IS RIGHT HERE, COME AND SIT IN THE CHAIR. He tells her 'No' but she tells him: YOU ARE NOT SAFE OUT ON YOUR OWN, and she takes his walking frame away and leads him back into his room, telling him: YOU ARE IN HOSPITAL, YOU NEED TO STAY SITTING IN YOUR CHAIR. Some time later, he is now in bed and appears to be asleep and there is now a one to one HCA sitting in a chair at the doorway watching him. [Site B day 3]

The typical technique when a person living with dementia was walking within the ward was to shadow them and to stand very close with arms folded. It was less common for the one to one carer to walk side-by-side with the person and their role was typically focussed on containing them at the bedside, directing them away from other bays, from talking to and shielding other patients, family or staff from them and diverting them away from the exits to the bay. As we can see during the care of this 79 year old man admitted with a fractured hip and described as 'confused', the main organisational force underlying and directing all of this work is the extreme fear of the patient in their care falling.

The room is darkened at the start of the night shift and the nurse and HCA are trying to get him back into bed: get back into bed. Patient: NO

The team respond: You will end up falling, You will end up falling, Get into bed it is night time. Patient: NO. The nurse: What will happen if you fall, If you fall you will be in hospital longer. The team get him back into bed and put the side rails up, but he is very agitated and pulling off his sheets. The HCA stands over him by his bed and as he repeatedly tried to get out of bed she keeps him within it: put your leg back in you will fall. [Site A day 18]

However, for families this was difficult, the impact of containment and restraint was that people became de-skilled and may lose their ability to walk. This has implications for outcomes and their discharge plans. Here, a daughter and son-in-law talk about her mother, who is 96 years old, has a diagnosis of dementia and was admitted with a hip fracture, I discuss with them that I had not seen her walk at all during her admission:

They won't let her walk, they are worried she will fall. She has walked with a zimmer for 7-8 years, so why cant she still use it? They say it's not safe. They say she has to pass the red thing first (the steady or rotunda) and once she can use that then she can move on to the zimmer. But she's too frightened of the red thing (the steady or rotunda) so she has never been able to try the zimmer and now she's lost the strength to walk, she cant get up herself now [...] They tried holding her up on either side, but she got a chest infection and didn't eat and then she was just not strong enough and they have not tried her since [...] They use the hoist (to get her from the bed to the chair or to the commode), but even we would find the hoist scary. How many times you say are you going to help her (We look at her legs- they are dark red/black and they tell me she has cellulitis). It is getting worse, her legs are going a funny colour and getting darker. [Site E day 7]

Dementia specialist teams were also available in some of the hospitals (Sites B and Site D). This was typically a small team (2 - 6 members of staff) who worked across a large number of wards and in some cases, all wards within the hospital, and thus they had limited time to spend within specific wards or individuals. Individual patients could also be identified by ward staff and medical teams as requiring support from the specialist team. These teams also only worked within 'office hours', 8:30 – 4:30, Monday to Friday. This group of dementia specialists were typically highly skilled HCAs, however, their time within wards and with individual patients was limited, and carried out in isolation to the ward teams. The ward and bay teams identified the patients who needed their support and expertise, but did not discuss strategies or approaches with them. In practice, their role was interpreted by ward staff as doing the interactional work with patients so that they did not need to do this. These specialists and their work with patients occurred in isolation and they described how they also felt unable to pass on their skills and knowledge of the person to other members of staff.

This outsourcing had wider impacts on the expertise of the ward team. It meant that the expertise of caring for a person living with dementia who was refusing care was never seen as the work of the ward, but the work of other people who could be temporarily brought into the ward. This also emphasised to the ward team that caring for large numbers of people with dementia was a temporary issue to be managed within their ward and not a permanent change in their patient population. The staff who had this one-to-one role typically had general HCA skills and only a small number talked about their expertise in dementia or had any continuity with the patient. The ward team and the individual typically identified the role of the one-to-one as containment and keeping the individual out of the way so that ward staff could continue uninterrupted with the timetable and schedule of the ward.

The impact of having a one-to-one care for the person with dementia was that the increasing level of surveillance, monitoring and the repeated requirements for them to remain at the bedside, and the repeated fears of falling and risk increased their anxieties of where they were and what was happening to them.

Visibility and invisibility

The work in the ward is driven by routine and timetables, but staff were also very responsive to individual buzzers, alarms and calls for immediate help. One impact of this was that patients who were silent or classified as 'sleepy' by ward staff were not a priority and so were less visible to staff during a shift, particularly if it was in the context of staff responding to multiple resistance within the bay and the wider ward. As a group, they were also unlikely to be assigned one-to-one care, with this assigned to patients identified as more actively agitated and resistant. We identified this pattern across all wards and sites (although there was a greater focus on this group within Site C). However, a silent or 'sleepy' patient or silence in a bay, does not mean all is well. Typically, within the ward, patients were lying in bed or sitting in their bedside chair and were often quiet or did not move, and appeared to be asleep. However, this does not mean they are not agitated or upset. In addition, there may still be body language that can be observed that gives some insight into their emotional state. Within encounters such as the one above (the 79 year old man admitted with a fractured hip and described as 'confused' at site A), the focus on patients who are actively resistant meant that the apparently silent 'sleepy' patient was not a priority. Within the same bay is a 84 year old man admitted with a hip fracture and pneumonia:

He is lying totally still and is slightly propped up in bed, his tiny head is lying back on the pillows, he has his glasses on and is staring up at the ceiling. He has an IV inserted in his left arm attached to a mobile drip stand -the night shift reported that he had pulled out the drip from his left arm in the night and they had to put it in again – I can see there is some spilt around his bed and there is a yellow 'caution wet floor' sign placed in the area around his bed. His bed covers are off and he is only partially covered in a thin sheet with his bare feet sticking out, which look very cold. He is also holding very firmly onto the side bars that are up on either side of his bed, but unlike previous days he is not pulling and rattling them or trying to get out of bed. He continues to lie very still in the bed for some time and then starts to fiddle with and then tries to pull the IV line out of his arm. However, it looks as though it has been very firmly and securely re-attached, there is lots of thick white bandages have been tightly wound around it covering a large part – over half - of his arm. He is unable to pull it out, but he keeps pulling. [Site A day 5]

Once a patient is made invisible by their condition, their quiet resistance became normalised for their admission, which could have significant impacts on care. As recognition of the patient as a 'person' declines, so to do their opportunities for rehabilitation and options for discharge.

Discussion

Our analysis has been drawn from detailed ethnographic observations that allowed the researchers to spend extended time within and across shifts observing specific bays and the individual staff and people living with dementia within them within hospital wards throughout England and Wales. This approach allowed us to identify the scale of resistance to care, but importantly, to explore the potential underlying reasons or triggers for a person's resistance to aspects of their care or admission. We have identified that these responses to the organisation and delivery of care at the bedside were typically rational to that patient's present ontology and perceptions. Importantly, we show that resistance typically did not take the form of isolated events where only one or a small number of individuals within the ward or bay resisted care. Every person we observed resisted care at some point during the observations. It was also typical for a number of patients to resist care at

the same time and within the same bay, particularly when staff delivered routine timetabled care seen by staff as essential care such as personal care, observation rounds, the medication rounds and mealtimes.

In isolation, each of these instances of resistance and refusal appeared relatively minor in terms of its overall impact on the ward, if not the patient, and often easily rectified. Tubes can be reinserted, machines reset, patients can be led back to bed, questions can be asked later. In combination, however, these minor acts of resistance, carried out by many patients, if not simultaneously, then in close association, had a significant influence on ward cultures, and approaches to and recognition of people living with dementia within the wards. Resistance to care has become something viewed by ward staff as a regular feature of a dementia diagnosis, and as both an expected and accepted feature of an individual with impaired capacity. This meant that staff respond to a person living with dementia who was calling out, shouting or walking in the corridor, not as a person expressing an underlying care need, but as a person demonstrating their diminished capacity, with this behaviour always attributed to their dementia diagnosis.

The more these acts of resistance occurred over a shift or a person's admission, the more staff normalised resistance, to the extent that it became interpreted as an everyday feature of ward life, a feature of a dementia diagnosis and who the person now is. The impact of this is that staff responses to people living with dementia become viewed as less urgent. An issue such as shouting or calling out becomes viewed by ward staff as a symptom of a pre-existing morbidity, rather than a feature of the acute admitting condition that the staff must quickly attend to. Rarely will the reason or intent of the patient's actions be discussed with the patient, beyond rhetorical platitudes such as 'where are you going?' and 'what's wrong darling?'. The heterogeneous 'dementia patient' mean that individuals become overshadowed by the perceived homogeneity of their condition.

The focus on completing the routines and 'rounds' of the ward timetables to schedule was always paramount, and dominated ward life. Ward teams approach to patient care was typically focused on carrying out very specific routine care for the person at the bedside, with a focus on completing the wider ward routines and timetables. This extended to their responses to resistance, which was typically to see these as potential interruptions to the work of the ward. and to carry on and try to complete that specific aspect of care. If a patient responded by resisting, or by saying 'no' to care, although staff would acknowledge this in their talk at the bedside, they typically reminded the person that they were in hospital and emphasised the rules of the ward. This highly repetitive talk was always directed at reminding the patient of their place in the world and the status of ward staff; they must all fit the expectations and timetables of the institution.

These exchanges provides ways of uncovering aspects of the loss of identity and social standing of people living with dementia during an acute admission. It is important to note how much of this talk was directed at trying to remind the patient of their place in the ward and the rules to be followed. Although staff gave the appearance of seeking permission and negotiating with the person, the delivery of care typically continued, with a tacit assumption of assent, focussed on obtaining the correct response from the person to allow care, which was already being carried out, to continue. However, this approach, which was highly repetitive, appeared to increase resistance, anxiety, and fear for people living with dementia.

The focus on the completion of ward routines and timetables meant that staff did not consider it possible to focus on what appeared to be low level resistance that was not an immediate priority or risk, particularly in the context of pressing demands of the ward routines of personal care, bed making, observation and medication rounds. It was often subtle signs that could be identified in a person's body language and changes in behaviour that indicated resistance or the potential for resistance. For example, when patients looked uncomfortable, displayed potentially defensive body

language (crossed arms), did not talk and were silent. Importantly, the challenge for ward teams in the context of busy wards was in recognizing these early signs, and feeling that they were able to respond to and prioritise them and that this constituted a valid form of work.

When ward routines and timetables were interrupted by a number of people resisting care, staff responses typically focused on short term management and containment so that they could focus on completion of the task in hand and the timetables of the ward, rather than responding to individual concerns. However, if these non-urgent signs of resistance were not acknowledged, then they typically escalated into the person developing a longer-term entrenched pattern of resistance. These perceived expectations of the timetables and their apparent conflict with patient needs could also create high levels of stress and anxiety for people living with dementia and also for ward staff.

The key ward staff response to people living with dementia within their care was one of containment and restraint. Although specific techniques had some variance between wards, the overall strategy was always the same: keeping the person living with dementia within their bed or sitting at the bedside. Across all sites, staff expressed high levels of anxiety about people leaving the bed or bedside, and this increased exponentially if they were walking in the bay, the wider ward and corridor or close to the ward entrance.

Multiple forms of containment and restraint were observed being used by all ward teams. Raising the side rails of the bed, or tucking bed sheets in tightly around the patient were both common and covert means to contain a patient within the bed. For those patients sitting at the bedside, the close placement of the mobile tray table, unreachable walking frames and technologies such as chair alarms were used to contain people to their chairs. Clinical technologies, including continence technologies (particularly full continence pads, bed pans, and commodes), 'specialing', medication and sedation, and secured medical equipment also limited or restrained movement from the bedside. Importantly, these approaches to patient care and their containment at the bedside was both a response to resistance, but were also frequently the trigger of resistance or cause of patient anxiety.

This contributes to what is seen commonly on wards throughout the acute hospital today, the outsourcing of dementia care. A key response to an identified pattern of resistance by people living with dementia within wards was to assign one-to-one agency HCA staff to care for them. This was a policy within almost all hospitals and wards as a way to support both people living with dementia and ward staff. In practice, this meant that once a person living with dementia was identified as resisting care within the ward, they could be classified as 'specialed,' a DoLs could be obtained and the ward could legitimately request additional support and assign an agency HCAs to provide one-to-one care to manage that person during their admission.

The role of the one-to-one carer was significant in that it was almost always interpreted as a role that required the restriction and containment of the person living with dementia to their bedside. To ensure that they fit within the rules of the ward and to limit their impact on the routines and timetabled schedules of care that dominated ward life. However, as well as having an impact on the person living with dementia, this approach could have powerful impacts on the wider ward culture. Ward staff appeared to be less concerned about de-escalating and supporting a person who was resisting care. This request could also function as a means to demonstrate to the wider hospital system that this was a ward and a team that was experiencing specific strains and required additional support staff. Caring for people living with dementia also came to be seen as not their core work of the ward, but to be outsourced. This in turn leads to the assignment of agency staff to such areas, contributing further to the stigma of care work for this patient group

Once a person is made invisible by their condition, their resistance became normalised as a feature of a dementia diagnosis, this could have significant impacts on care. Such processes, generated from the restricted repertoire of work possible within the routines of these settings, leads to cultures of dehumanisation for patients, but also has negative impacts on staff, who do not consciously create these cultures but operate within them, leading to potential emotional burnout and exhaustion. It was not uncommon for staff to want to avoid assignment to bays or areas of wards or units that admit high numbers of people with dementia. The work in wards is driven by routine and timetables, but staff were also very responsive to individual buzzers and calls for immediate help. One impact of this was that patients who were silent or classified as 'sleepy' by ward staff were not a priority and became less visible to staff during a shift, particularly if it was in the context of staff responding to multiple resistance within the bay and the wider ward. As a group, these people living with dementia were also unlikely to be assigned one-to-one care, less likely to get staff attention that those patients identified as more actively agitated and resistant. However, as recognition of the patient, as a person, declines, so to do their opportunities for rehabilitation and options for discharge.

Conclusion: interventions and change

In response, we believe that our ethnographic 'thick description' is particularly applicable for developing organisational and interactional training and interventions at ward level. Ethnography provides ways to connect ward staff with key issues by providing detailed 'real' empirical examples178 of care that bring these issues alive and provoke better awareness. We believe (and have found in our discussions with ward staff) that these detailed accounts of the organization and delivery of care within wards will support and encourage staff to develop a strong connection and understandings of the social world of their ward, the ways in which the organization of work influences them, how ward cultures of care develop and become normalized, and the perspectives of people living with dementia and their families. We believe that it is only through an empirical detailed understanding of the complex social relationships involved in the current provision of care for people living with dementia in acute wards and the significant impacts this has on them, their families and carers, and for ward staff, can we identify ways in which to deliver high quality care for people living with dementia that also supports ward staff.

Summary

We have provided a detailed analysis of the experiences of care and the impacts of the culture and organization of the wards and interactional work of delivering care at the bedside from the perspectives of people living with dementia, their family carers, and ward staff. We have identified the ways in which resistance to care manifests, is recorgnised by staff and the routine responses to it. We have also examined the impacts of these responses over time. A number of studies have highlighted the difficulties of caring for people living with dementia within the acute hospital setting (Tadd et al 2011, Dewing and Dijk 2014), however, there is an evidence vacuum in understanding how their care can be improved within the acute setting (Royal College of Psychiatrists, 2013), and few studies have demonstrated how to practically address these challenges. In response, within this chapter, we demonstrate how we are using our findings to develop training and test the feasibility of interventions to improve the quality and humanity of care for this key population and for the ward staff caring for them.

Within this final chapter we focus on our approaches and strategies to build upon and utilise the study findings to (1) consult with, raise awareness and inform wider user communities and publics about our findings and the experiences of care in acute wards for people living with dementia, their family carers and for ward staff; (2) extend dissemination and delivery of training to nurses and HCAs working in acute wards via open access on-line training; and (3) develop and test interventions at ward level to improve the quality and humanity of care people living with dementia receive during a hospital admission. To do this, we have:

- Involved people living with dementia, their carers, families, and the wider publics, in the research process and to discuss initial findings, via an ongoing dissemination and consultation process in collaboration with the arts by developing a programme of dementia friendly film screenings and festivals.
- Presented the initial findings of our research to nurses, students, care staff, patients, advocacy groups and experts from a variety of disciplines in the UK and Europe.
- Worked with the arts to support the integration of people living with dementia and other potentially excluded audiences to participate in both the arts and the wider experience of being part of cultural and artistic venues. We have also pioneered a toolkit (in collaboration with the BFI Film Hub Wales) and supported staff training to facilitate this.
- To extend and enhance our community of interest we have developed social media programmes to extend the reach of our research via a coordinated cross-platform campaign of film content, training and research updates via open access social media platforms (dedicated website, twitter, facebook, Instagram, youtube).
- Established our brand 'storiesofdementia' via accessible logos and illustrations to create a distinct, memorable and consistent identity that reflects the overall objectives and values that are at the heart of our research and to ensure that our outputs instantly recognisable as belonging to our project.
- Collaborated with Dementia UK and Admiral nurses on developing:
 - o no-cost interventions or 'hacks' for acute wards
 - o short training (n=17) and 'trigger' (n=12) films for nurses and HCAs working in acute wards available via open access on-line training.
- Conducted a feasibility study within one 'laboratory' ward to introduce three interventions or 'hacks' and to establish the feasibility of introducing them in acute wards.

Our detailed analysis has identified features of the social and organisational context of frontline care and key moments within the hospital system and ward routines and timetables which we believe are pivotal opportunities for change. In response, we have developed small changes or 'hacks', which can be made that do not require 'permission', cost money or demand involvement of the hospital management or wider governance structures. We believe that these changes can improve both the quality and humanity of care for people living with dementia and also improve the working lives of ward staff. An individual nurse or HCA may believe they lack the power or influence to make large transformations to their hospital, but small changes can have significant impact on the local culture of the ward, the ward team and on patient care.

Our expertise is becoming sought after and we are extending our training and interventions to involve other hospital Trusts and other areas of care (delirium) that impact on people living with dementia during an acute hospital stay. The findings have also informed a further study currently underway: Understanding how to facilitate continence for people living with dementia in acute hospital settings: raising awareness and improving care (NIHR HS&DR: 15/136/67).

Consultation, raising awareness and informing wider user communities and publics about our findings

Importantly, this project takes a creative approach to reducing social inequalities experienced by those affected by dementia, improving care for people living with dementia, and disseminating, engaging and consulting with people about our research through a sustained collaboration with the arts. Although the focus of our work is improving care in acute settings, we know that the experiences of people living with dementia within that setting is a reflection of the stigma and invisibility people experience more widely within society. In response, we are working with the arts to support the integration of a potentially excluded audience, providing the opportunity for them to participate in the general programme and the wider venue experience.

Collaboration with the arts

We are working in partnership with key stakeholders to develop a community of interest to promote dissemination of key findings and debate about what care for people living with dementia should look like, with people living with dementia, their carers and families, and wider publics. This builds on a long-term collaboration with Chapter (National Cinema for Wales http://www.chapter.org/), and the British Film Institute funded Film hub Wales (http://www.chapter.org/welcome-film-hub-wales). Specifically:

Dementia Friendly Film Screenings:

In collaboration with Chapter (https://www.chapter.org) and the BFI Film hub Wales (https://filmhubwales.org), this project takes a creative approach to reducing social exclusion in dementia by delivering a monthly programme of 'dementia friendly' and 'supportive' film screenings for people living with dementia and their amily carers. Running since 2016, in 2017 we held 20 'dementia friendly' screenings from classic musicals (including Grease Singalong and Calamity Jane) to new release British films (Eddie the eagle, A Cat Named Bob and Viceroys House), general release (La La Land) and independents (The Straight Story), attended by 727 people.

This collaboration is developing the evidence base for a sustainable model of enjoyable and inclusive film events that encourage and enable people living with dementia, their carers and wider film audiences to come together to share their love of film. Working with the arts to support the integration of potentially excluded audiences to participate in the arts and the wider experience of being part of cultural and artistic venues. Importantly, this also grows a direct community for dissemination and engagement with the research. This collaboration also provides access to partners across Wales, UK, and Internationally via the BFI.

Training

In collaboration with Film hub Wales (http://filmhubwales.org) we are working to improve access to the arts and increase diversity in cinema audiences via the development of toolkits to support venues across the UK to increase access for people living with dementia and other potentially excluded audiences (http://filmhubwales.org/resources/access-all). So far, four Arts Centres and Independent cinemas are using our model to launch Dementia Friendly Film Screenings across Wales.

- Dementia friendly training for Cinema Staff (South Wales): 100 participants attending Dementia Friends Sessions.
- 'Opening Doors' Diversity training days: these training days (Film Hub Wales, Chapter, BFI), are designed to support film venues across Wales in diversifying their audiences. They aim to welcome community groups who do not currently use their venue. These training days now include a session on creating dementia friendly communities and spaces.

Tinted lens curated film season

Tinted Lens: a programme of curated film events that explores the ideas found within new-release, cult and classic film, with a focus on understandings of the mind, human behaviour, memory, the life-course, and ageing. This programme of monthly events provides an ongoing forum for dissemination, engagement and debates about ongoing research. The monthly events over 3 years (2015- present) have included: Still Alice, Robot and Frank, Solaris, The Possibilities are Endless, The Falling, Ida, and Mr Holmes, The Lost Weekend, Irrational Man, The Lobster and The Danish Girl. These have an average audience of 40-50 members of the public attending both the film and the discussion (total: >1,200 participants).

Tinted Lens Festival of the mind, memory and ageing

In collaboration with Chapter and the Film hub Wales (BFI) we held a festival to celebrate Dementia Awareness Week (May, 2017). We believe this was the first of its kind, and our research and key study findings reached over 500 people, including people living with dementia, their carers and families, government ministers, policymakers, practitioners, dementia charities, and third sector organisations. The festival included film screenings (including dementia friendly screenings and public debates); interactive research projects, research stalls and research presentations and posters; performances and workshops from a range of theatre company (Everyman and Re-Live), arts and craft interactive workshops; and dementia friendly choirs (Forget-me-not chorus). Evaluation is in process and outputs will include a short film. The success of the festival has meant that this is now an annual event, with the next planned for May 2018.

Dissemination to ward staff

We have begun the process of disseminating the findings of our research in an effort to improve the care of people living with dementia within the acute setting. In order to best reach as many nurses and HCAs as possible, we have provided multiple pathways of engagement, from in situ ward 'hacks' to online open access films.

Need for training

The majority of nursing staff (89%) working in the acute setting have identified working with people living with dementia as challenging (Alzheimer's Society 2008), with healthcare professionals within the acute setting identified as lacking the necessary skills and knowledge to care for this patient population (Scottish Government, 2010; Department of Health, 2009; Tadd et al, 2011). Yet without the appropriate training and support, there is a recognition that healthcare staff will become resentful, demoralised and cut themselves off from patients, all things that can lead to the depersonalisation and dehumanization of people in their care (Cornwell, 2012). With the failure to provide appropriate training for hospital staff in caring for people living with dementia identified as a key contribution to their poor outcomes and long inpatient stays (Alzheimer's Society, 2007). Training in the care and support of people living with dementia must also be part of the continuous professional development for nurses (Department of Health, 2009), with a further emphasis on the need for training all staff working in health or social care settings (Department of Health and Social care, 2011; Alzheimer's Society, 2013). Training is recognized as the key to reducing stigma (Department of Health, 2009) and delivering dignity in care (Department of Health (2015).

From 'bundles of care' to 'hacks'

We initially used our findings to develop a number of 'bundles of care'. The current care bundle for dementia patients in hospital advocates a series of measures such as using a reminiscence artefacts, developing communication strategies and having a communal dining room where possible (Brooker et al 2013). Clearly such measures are inappropriate within many acute areas where patient turnover is high (such as MAUs), and where medical interventions are often complex and have priority, and staff are focussed on clinical outcomes. Indeed, when observing staff in their day-to-day interactions with patients living with dementia, the main issues they raised was the need for support and training on how to actually care for people living with dementia in their wards and how to respond to and manage resistance to care when they are caring for people at the bedside.

In response, we have used our findings to develop a number of 'bundles of care', which each comprises of a small, straightforward set of five evidence-based practices. We have piloted these bundles within undergraduate nursing modules and consulted with ward staff. Our goal was to

provide structured ways to improve the processes of care, interactions with patients, and inform ward culture, to support people with dementia and ward staff:

- · Caring for people with dementia in acute wards,
- Caring for people with dementia in MAU,
- Developing ward cultures that supports people with dementia,
- Caring for people with dementia who are medically fit to leave.

However, following a consultation process with ward staff in a number of hospital Trusts and Admiral nurses, we identified that there was already 'bundle fatigue' (meeting with nurses and ward managers, site A, 2017) amongst nurses working in acute wards. The key finding of our consultation was that ward staff were wary and cynical of any intervention that could involve new paperwork or additional forms entering their ward because they were always viewed as adding to their workload. Instead, they asked for simple no-cost interventions, techniques, and training, that they could implement within their practice and their wards without having to seek 'permission' (meeting with Admiral Nurses, 2017) from the wider hospital administrative and executive systems. As one nurse reported 'how can we make changes when I still can't get permission to put a nail in the wall to put a clock up in the ward!' (meeting with ward staff, site A, 2017).

In response, we held a national expert 'hackathon' event (1st to 2nd November 2017, Birmingham) with Dementia specialist nurses and Admiral Nurses invited from across England and Wales, specifically to refine our approaches to implementation. We focussed on exploring the ways in which our study findings could be adapted into implementable and maintainable solutions on acute wards, in the form of small 'hacks'. We use the 'hack' analogy to describe a particular problemsolving approach, or 'social hacking' focussed on identifying ways to solve social issues. We believe this approach can be applied to acute care. We are using our analysis to identify key moments within the hospital system and ward routines and timetables which we believe are pivotal opportunities for change. From this, we have then identified a number of simple 'hacks' that allow us to intervene in ways that we believe could change ingrained behaviour and improve care. These are simple no-cost interventions that we believe can be engrained into everyday practice. These hacks are designed to improve the experience of patients living with dementia, but also the experience of ward staff caring for them during their admission to an acute hospital setting. A key aspect of this process is to continually test and refine the 'hacks' to learn from them and to improve them.

The hackathon event identified three 'hacks', each covering a specific area of everyday care that we believe are key to improving care for people living with dementia: (1) communication, (2) mealtimes and hydration, and (3) movement and rehabilitation within the ward. These are being used to:

- inform on-line training films available to hospital staff (open access via our blog).
- Inform the development of a masterclass and future MOOC (NIHR funded).
- Inform the design of an NIHR funded feasibility study within one acute 'laboratory' ward to test and refine the 'hacks' and to establish their feasibility within the acute ward setting.
- Provide the foundations for a collaborative funding application for a multi-centre intervention study.

On-line training in collaboration with Dementia UK

Our development work with Admiral nurses and Dementia UK will form the basis of our training films, masterclasses and MOOC for nurses, HCAs and ward staff in caring for people living with dementia within acute ward settings. It will also be integrated into Cardiff University undergraduate and postgraduate nursing training (via Cardiff University) and also available via our own dedicated website (www.storiesofdementia.com). So far, we have published 29 short training films that include:

- Training films that include 'Top tips', 'Do', 'Don't', 'If I could change one thing', and 'My biggest challenge' (n=17)
- Awareness raising stories and 'trigger' films (n=12) from the perspectives of people living with dementia, carers and families, and nurses.

Carers perspectives

Our ongoing work with the Carer Steering Group is focused on developing ways in which the findings generated by the study should be communicated to user communities. So far, in collaboration we have developed:

- Short films with people living with dementia and their family carers discussing their experiences of hospital care to raise awareness and to be used as 'trigger' films in wards to support ward change and development to improve care for people living with dementia.
- A short report and factsheet for carers and families focusing on 'top tips' and things that are important for families to be aware of and to consider during a hospital admission for a person living with dementia.

The Carers Steering Group have provided invaluable input towards making sure the language used in the reports and factsheets is both appropriate and accessible.

NHS Trusts

We have presented our findings at a number of NHS sites and hospital level meetings including Dementia Boards, Nurse Divisional meetings, Senior Nurse meetings, research days and working groups. These meetings include managers of acute settings and senior nursing teams. We are currently working with nurses and acute wards across a number of hospital Trusts, teams of dementia specialist nurses, and Admiral Nurses across England and Wales. We plan to involve these teams and sites in our ongoing work to develop and implement interventions and 'hacks' at ward level.

Developing no-cost interventions at ward level

We are currently in the process of conducting a feasibility study that will use the study findings within one acute 'laboratory' ward to develop innovative and practical interactional and organisational interventions or 'hacks' to improve the quality and humanity of care people living with dementia receive during a hospital admission. We are assessing the feasibility of implementing them at little or no cost in the clinical area.

Theory

The conceptual framework guiding this feasibility study is adapted from Deming's theory of profound knowledge, which is a management philosophy grounded in systems theory (known as the system of profound knowledge; SoPK) (Deming, 1993). It is based on the principle that each organization is composed of a system of interrelated processes and people which make up system's components. The success of all workers within the system is dependent on management's capability to orchestrate the delicate balance of each component for optimization of the entire system. Whilst originally intended as quality tool in manufacturing businesses its focus on leadership means that it has been adopted across a range of healthcare settings. Deming believed that profound knowledge generally comes from outside the system and is only useful if it is invited and received with an eagerness to learn and improve. A system cannot understand itself without help from outside the system, because prior experiences will bias objectivity, preventing critical analysis of the organisation. Critical self-examination is difficult without impartial analysis from

outside the organisation and so the purpose of the study is thus: to assist nurses in adopting small changes to make large differences.

This journey, however, should not be under-estimated, because it requires leadership from within and a desire to obtain and us (profound) knowledge to guide this process. Profound Knowledge is made up of four interrelated components: Appreciation of a system; Theory of knowledge; The psychology of change; Knowledge about variation. These four components cannot be separated and a knowledge of psychology, variation, theory of knowledge, and appreciating the processes of a system, must be managed as a delicate balance; they make up systems.

Feasibility study: an experienced-based co-design framework

In response, the feasibility study uses an experienced-based co-design framework (Robert et al, 2013) as described by Catherine Dale (Kings Fund, 2008), which involves gathering experiences from patients and staff through in-depth interviewing, observations and group discussions, identifying key 'touch points' (emotionally significant points) and assigning positive or negative feelings. The approach was designed for and within the NHS to develop simple solutions that offer users a better experience of treatment and care. This approach has already been used in a range of clinical services, including cancer, diabetes, drug and alcohol treatment, emergency services, genetics, inpatient units, intensive care, mental health, orthopaedics, palliative care and surgical units. However, it is relatively untested in the field of dementia care and, in particular, no work has focussed on people living with dementia receiving acute care for unrelated health problems in non-specialist settings.

To build upon and utilise the study findings to inform ward care, we held a 'Hackathon' event (1/2nd November 2017 in Birmingham) that allowed for input, discussion and feedback from representatives of Dementia UK and a group of specialist and Admiral nurses with experience of caring for people living with dementia in the acute setting. This event produced a lively debate and discussion across two days and generated three 'hacks', which were identified as both useful and appropriate for feasibility testing within an acute care setting.

Feasibility study: Implementation phase

This implementation phase is a feasibility study being carried out in collaboration with Aneurin Bevan University Health Board (ABUHB), South Wales. We have approvals in place and are carrying out this feasibility study within a large general hospital, the Royal Gwent Hospital, Newport, and to work in collaboration with the deputy ward sister of the Acute Medical Frailty Unit (D2East). Professor Sue Bale (Director of R&D, ABUHB), has also take day-to-day responsibility for the project locally, including the identification and recruitment of participants. The implementation of the feasibility study is being carried out by team members (Featherstone and Harden) Both have significant experience of working in collaboration with nursing staff within acute hospital wards, and of conducting research that includes people living with dementia.

We have introduced the 'trigger' films and potential 'hacks' and discussed them with frontline nursing and HCAs staff to support discussion about the challenges of caring for people living with dementia within their ward and how to implement the ideas and possible solutions via the 'hacks' produced by the forum. These 'hacks' were all identified as potentially useful by ward staff, are currently being implemented and evaluated in turn, to test the feasibility of implementing them in an acute ward and to provide an indication of their effectiveness.

Research aims and objectives

This study explores the feasibility of introducing and implementing the low to zero cost recommendations of the MemoryCare project within the acute care setting. Our key research questiona are: Is introducing and implementing the recommendations of the MemoryCare project or 'hacks' within the acute care setting feasible? Can they improve the delivery and experiences of care for people living with dementia, for both staff and patients?

The objectives are to examine and assess if the findings of our detailed ethnographic analysis can be implemented and if they can improve the delivery of care for people living with dementia, for both staff and patients. Within one acute ward the feasibility study is:

- Providing ward staff with the opportunity to discuss the three 'hacks' identified from analysis of the primary study, and the discussion of its finding with practitioners and experts at the 'Hackathon' event (held 1/11/2017).
- Assess the potential usage of the three 'hacks' with ward staff for improving care for people living with dementia, and the feasibility of implementing each of the three 'hacks' within their ward.
- Run co-design group with ABUHB ward staff and users over a four month period to work on agreed improvements in the form of 'hacks' or 'hacks'
- Evaluate using patient outcome measurements and patient and staff satisfaction surveys to establish the feasibility of using our 'hacks' to inform quality improvement at ward level.
- Further refine 'hacks', and use this to inform a larger-scale acute-setting wide pilot to examine whether the interventions and approach to supporting change at ward level is scalable.

Implementation and evaluation

We follow the process evaluation guidelines of complex interventions as prescribed by the Medical Research Council (see Moore et al 2015). The rationale for using the MRC process evaluation model is that it aims to provide a more detailed understanding which is needed to inform policy and practice and may be conducted within feasibility testing phases, alongside evaluations of effectiveness or alongside post-evaluation scale-up. We use this approach to provide both descriptive quantitative information on fidelity (quality), 'dose' (completeness/satisfaction), and reach (participation rate). Qualitative data is being collected through iterative interviews with nurses and carers to generate themes which will then be used to explain and frame the quantitative findings. This is achieved through examining aspects such as:

Implementation: In this context of this study, the term 'implementation' describes both the development – evaluation – implementation process, and the implementation delivery during the evaluation period. We are carrying out observations of nurses and Health Care Assistants (HCAs) within the ward, which will concentrate on the everyday work of nurses, Health Care Assistants and other ward staff as they are involved in implementing the 'hack' or 'hacks'. This involves 15 days/shifts of observation, focussing on the structures, resources and processes through which delivery is achieved, and the quantity and quality of what is delivered.

Mechanisms of impact: We are also interested in examining how the intervention activities, and participants' interactions with them, trigger change. In addition to the observations, we are also (where possible) carrying out short ethnographic (within observation) interviews with ward staff (n=30).

Context: External factors and their influence on the delivery and functioning of interventions within the ward are also key and in response, we will also evaluate the feasibility of using the interventions and their effectiveness using data from feedback forms, provided to both staff and patients (and/or their families/carers).

Process evaluation involves critically observing the work of intervention staff, in this case nurses and HCAs, and it is important that the research team sustain good working relationships, whilst remaining sufficiently independent for evaluation to remain credible. This has been facilitated by the relationship already developed in the initial NIHR study as the site being used now was also the original pilot site.

Using the MRC process evaluation model allows us to:

- provide a clear description of the intended intervention,
- observe how it was implemented and
- how (if) it worked.

The findings of this feasibility study will be used to, where necessary, refine our hacks, and we hope to use this to inform a larger-scale pilot to examine whether the 'hacks' and approach to supporting change at ward level is scalable.

Acknowledgements

This project was funded by the National Institute for Health Research NIHR HS&DR researcher led funding stream: MemoryCare: The management of refusal of food, drink and medications by people with dementia admitted to hospital with an acute condition (project number 13/10/80). The views and opinions expressed therein are those of the authors and do not necessarily reflect those of the HS&DR researcher led funding stream, NIHR, NHS or the Department of Health. The full version of this report will be published within the National Institute for Health Research Journals Library.

We could like to thank our project and carers steering group for all their support and for keeping us focussed on our goal of improving the quality and humanity of care people living with dementia receive during an acute hospital admission. We particularly thank Rosie Tope, Jackie Askey, Anne Davies, Peggy Martin, Viv Morgan, Julie Stacey, Betty Roderick, David Jones, Lynne Jeffrey, Chris Jones, and all the members of the Carers support network.

We are of course, extremely grateful to all the people living with dementia, their families and carers, and the nurses and healthcare assistants that have shared their stories, supported our work, and given us the encouragement to use our work to improve care for people living with dementia, and also to improve the working lives of nurses and healthcare assistants working in the acute setting. We particularly thank Gaye Birch, the first carer who agreed to be involved in our research during her husband's hospital admission and who has been a great and continued supporter of the team and our research since then.

We also thank all the staff within Anaerun Bevan University Health Board, particularly Professor Sue Bale and the Research and Development team, the Dementia Board, Brognah Scott (Director of Nursing), Lin Slater (Deputy Director of Nursing), Lorraine Edmunds (Dementia Lead) and all the nurses working with her.

We have drawn on a wide range of expertise in delivering this project and our wider programme of communicating our work: Nick Daw (Director of Photography http://nickdaw.com), Tyler Chambers (Social Media), Sophie Nightingale (Branding, web design and illustration https://www.sophienightingale.net), Chris Hartwill (Director https://www.rsafilms.com/uk/directors/film-and-tv-directors/chris-hartwell/featured/), and Kelly Hewitt (Producer, http://www.callboxdiary.com/callbox/kelly-hewitt/).

We are also indebted to the teams at Chapter (<u>www.chapter.org</u>) and Film Hub Wales (<u>www.filmhubwales.org</u>), who all have a deep commitment to increasing access to the arts and

using film to reduce social isolation and stigma. They have always been enthusiastic about bcollaborating with us to improve access to the arts and film for people living with dementia and their families and we look forward to continuing this work: Sally Griffith (Director of Film & Cinema, Chapter and Director of Film Hub Wales), Claire Vaughan (Programme & Learning Officer, Film & Cinema) Ellie Russell (Dementia Project Officer), Hana Lewis (Strategic Hub manager) and Lisa Nesbitt (Development Officer) and Toki Allinson (Film Audience Network Project Officer).

References:

Age UK, 2012 Deliving Diginity: Securing dignity in care for older people in hospitals: A report by the Commission on Dignity in Care Delivering Dignity @ http://www.nhsconfed.org/-/media/Confederation/Files/Publications/Documents/Delivering_Dignity_final_report150612.pdf accessed 12/02/18

Allen, D. A., & Lyne, P. A., 1998. Nurses' flexible working practices: some ethnographic insights into clinical effectiveness. *Clinical Effectiveness in Nursing*, 1(3), 131-138.

Alzheimer's Society, 2007. Annual Review. London: Alzheimers Society.

Alzheimer's Society, 2008. Annual Review. London: Alzheimers Society.

Alzheimer's Society, 2015. *Dementia 2015: Aiming higher to transform lives.* London: Alzheimers Society.

Alzheimer's Society, 2016. Annual Review. London: Alzheimers Society.

Alzheimer's Society, 2016. Fix Dementia Care: Hospitals London: Alzheimer's Society

Alzheimer's Society, 2017. Annual Review. London: Alzheimers Society.

Andrews, J., Fellow, R.C.N. and Butler, M., 2014. *Trusted to Care: An independent Review of the Princess of Wales Hospital and Neath Port Talbot Hospital at Abertawe Bro Morgannwg University Health Board*. Welsh Government.

Atkinson, P., Delamont, S. and Housley, W., 2008. *Contours of culture: Complex ethnography and the ethnography of complexity*. Rowman Altamira.

Baikie, E. (2002). The impact of dementia on marital relationships. *Sexual and Relationship Therapy*, 17(3), 289-299.

Bamford, C., Lamont, S., Eccles, M., Robinson, L., May, C. and Bond, J., 2004. Disclosing a diagnosis of dementia: a systematic review. *International journal of geriatric psychiatry*, 19(2), pp.151-169.

Bamford, S., 2011. Women and dementia—not forgotten. Report by the UK International Longevity Centre.

Banicek J., 2010. How to ensure acute pain in older people is appropriately assessed and managed. *Nursing Times*; 106: 29, 14–17.

Beerens, H.C., 2013. "Factors associated with quality of life of people with dementia in long-term care facilities: A systematic review." *International Journal of Nursing Studies*.

Black. N., 1994. Why we need qualitative research. J Epidemiol Commun Health 48:425-6

Blumer, H., 1954. What is wrong with social theory?. American sociological review, 19(1), pp.3-10.

Bonda, M.J., Clark, M.S, Davies, S. 2003. The quality of life of spouse dementia caregivers: changes associated with yielding to formal care and widowhood *Social Science & Medicine* 57 2385–2395

Borbasi, Sally, et al. "Health professionals' perspectives of providing care to people with dementia in the acute setting: Toward better practice." *Geriatric Nursing* 27.5 (2006): 300-308.

Bradley E., Webster T., Baker D., Schlesinger M. & Inouye S. (2005) After adoption: sustaining the innovation. A case study of disseminating the Hospital Elder Life Program. Journal of the American Geriatrics Society 53, 1455–1461.

Brauner, D., 2009. Adherence to medication in patients with dementia: problems and solutions. *Geriatrics & Aging*, 12(5), pp.259-263.

Bridges, J., et al., 2013. Capacity for care: meta-ethnography of acute care nurses' experiences of the nurse-patient relationship. *Journal of Advanced Nursing*, 69(4): p. 760-772.

Brooker, D., Leung, D., Bowley, K., Etches, C., Bray, J., Smith, P. and Upton, D., 2013. The Dementia Care Bundle. Improving the Quality and Safety of Hospital Care for Patients with Acute Physical Illness Who have Co-existing Dementia. The Health Foundation. Abgerufen von http://patientsafety. health. org. uk/sites/default/files/resources/the_dementia_care_bundle. pdf.

Caddell, L.S., Clare, L., 2010. "The impact of dementia on self and identity: A systematic review." *Clinical Psychology Review* 30.1: 113-126.

Caracelli, V. J. 2006. Enhancing the policy process through the use of ethnography and other study frameworks: A mixed-method strategy. *Research in the Schools*, 13(1), 84-92.

Care Quality Commission., 2011 *Dignity and Nutrition Inspection Program National Report* London: DoH

Care Quality Commission, 2013. The state of health care and adult social care in England in 2013/14(Vol. 763). The Stationery Office.

Care Quality Commission, 2014. The state of health care and adult social care in England in 2013/14(Vol. 763). London: The Stationery Office.

Care Quality Commission, 2017. The state of health care and adult social care in England 2016/17 London: The Stationery Office.

Charmaz, K., & Mitchell, R. G. 2001. Grounded theory in ethnography. Handbook of ethnography, 160-174.

Chenoweth, L., et al., 2101 "A systematic review of what factors attract and retain nurses in aged and dementia care." *Journal of Clinical Nursing* 19.1-2: 156-167.

Cheong CY, Tan JAQ, Foong Y, Koh HM, Chen DZY, Tan JJC, Ng CJ, Yap P. 2016. 'Creative Music Therapy in an Acute Care Setting for Older Patients with Delirium and Dementia' *Dementia and Geriatric Cognitive Disorders* 6 268-275

Coffey, A. and Atkinson, P., 1996. *Making sense of qualitative data: complementary research strategies*. Sage Publications, Inc.

Collins, N., Blanchard, M.R., Tookman, A. and Sampson, E.L., 2010. Detection of delirium in the acute hospital. *Age and ageing*, 39(1), pp.131-135.

Cooper, C., et al. "A systematic review and meta-analysis of ethnic differences in use of dementia treatment, care, and research." *American Journal of Geriatric Psych* 18.3 (2010): 193-203.

Corbin, J., Strauss, A.,1990., Grounded theory research: procedures, canons and evaluative criteria

Cornwell, J., 2012. The care of frail older people with complex needs: time for a revolution. London: Kings Fund

Creditor, M. C., 1993. Hazards of hospitalization of the elderly. *Annals of internal medicine*, 118(3), 219-223.

Curtis, S., Gesler, W., Smith, G. and Washburn, S., 2000. Approaches to sampling and case selection in qualitative research: examples in the geography of health. *Social Science & Medicine*, 50(7), pp.1001-1014.

Cutson T, Ramos K, Grant J, Moore M, Fulton J. 2016. 'Swinging for the fences: Managing challenging behaviours at end of life' *Journal of Pain and Symptom Management* 51(2) 366

Davies, S., Nolan, M., Brown, J., Wilson, F., 1999. *Dignity on the ward: promoting excellence in care*. Help the Aged, London.

Dawson, P. and Reid, D.W., 1987. Behavioral dimensions of patients at risk of wandering. *The Gerontologist*, 27(1), pp.104-107.

Daykin, N., & Clarke, B. (2000). 'They'll still get the bodily care'. Discourses of care and relationships between nurses and health care assistants in the NHS. *Sociology of Health & Illness*, 22(3), 349-363.

De Boer, M. E., Hertogh, C. M., Dröes, R. M., Riphagen, I. I., Jonker, C., & Eefsting, J. A. (2007). Suffering from dementia—the patient's perspective: A review of the literature. *International Psychogeriatrics*, 19(06), 1021-1039.

Dementia Action Alliance (2012) Dementia Friendly Hospital Charter https://www.dementiaaction.org.uk/assets/0001/8146/DAA_Dementia_Friendly_Hospital_Chart er_Booklet_06-2015.pdf accessed 12/02/2018

Deming, W.E., 1993. A system of profound knowledge.

Department of Health., 2009. Living Well With Dementia: a national dementia strategy London:

DoH

Department of Health and Social care, 2011. *Common Core Principles for Supporting People with Dementia. A guide to training the social care and health workforce* London: DoH

Department of Health, 2013. Dementia: A state of the nation report on dementia care and support in England London: DoH

Department of Health, 2015. Prime Minister's Challenge on Dementia 2020.

Dewing, J., Dijk, S., 2014. What is the current state of care for older people with dementia in general hospitals? A literature review. *Dementia* 15(1) 106-124

Dixon-Woods, M. (2003). What can ethnography do for quality and safety in health care?. *Quality* and Safety in Health Care, 12(5), 326-327.

Eggenberger, E., Heimerl. K., Bennett. M.I., 2012 "Communication skills training in dementia care: a systematic review of effectiveness, training content, and didactic methods in different care settings." *International Psychogeriatrics* 1.1: 1-14.

Elliott, K.J., et al. "Building capacity and resilience in the dementia care workforce: a systematic review of interventions targeting worker and organizational outcomes." *International Psychogeriatrics* 24.6 (2012): 882.

Emerson, R. M., Fretz, R. I., & Shaw, L. L. (2011). *Writing ethnographic fieldnotes*. University of Chicago Press.

Francis, R., 2013. Report of the Mid Staffordshire NHS Foundation Trust Public Inquiry: Executive Summary Vol. 947. London: The Stationery Office

George, J., Long, S., Vincent, C., 2013. How can we keep patients with dementia safe in our acute hospitals? A review of challenges and solutions. *Journal of the Royal Society of Medicine*, 106(9), 355-361.

Glaser, B. and Strauss, A., 1967. *The discovery of grounded theory*. London: Weidenfeld and Nicholson, 24(25), pp.288-304.

Goffman, E., 1959. The Presentation of Self in. Butler, Bodies that Matter.

Goldberg, S. E., Whittamore, K. H., Harwood, R. H., Bradshaw, L. E., Gladman, J. R., & Jones, R. G., 2012. The prevalence of mental health problems among older adults admitted as an emergency to a general hospital. *Age and ageing*, 41(1), 80-86.

Goodson, L., Vassar, M. 2011. An overview of ethnography in healthcare and medical education research *J Educ Eval Health Prof*, 8: 4

Graham, N., Lindesay, J., Katona, C., Bertolote, J. M., Camus, V., Copeland, J. R., Wancata, J., 2003. Reducing stigma and discrimination against older people with mental disorders: a technical consensus statement. *International Journal of Geriatric Psychiatry*, 18(8), 670-678.

Green, J. 1998. Commentary: grounded theory and the constant comparative method. *BMJ* (*Clinical research ed.*), 316(7137), 1064.

Greenhalgh, T., Swinglehurst, D., 2011. Studying technology use as social practice: the untapped potential of ethnography. *BMC medicine*, 9(1), 45.

Guba, E.G. and Lincoln, Y.S., 1994. Competing paradigms in qualitative research. *Handbook of qualitative research*, 2(163-194), p.105.

Hammersley, M., 1987. What's Wrong with Ethnography? Methodological Explorations. London: Routledge

Hammersley, M., 2004. Action research: a contradiction in terms?. *Oxford Review of Education*, 30(2), pp.165-181.

Hammersley, M. 2006. Ethnography: Problems and prospects. *Ethnography and Education*, 1(1), 3-14.

Hammersley, M., Atkinson, P., 1989. Ethnography: Principles in practice. London: Routledge.

Harris, P. B. (2012). Maintaining friendships in early stage dementia: Factors to consider. *Dementia*, 11(3), 305-314.

Hart B.D., Birkas J., Lachmann M., Saunders L. 2002. Promoting positive outcomes for elderly persons in the hospital: prevention and risk factor modification. *American Association of Critical-Care Nurses: Clinical Issues* 13(1) 22–33.

Health Foundation., 2011. *Spotlight on dementia care: A Health Foundation improvement report*. London: The Health Foundation.

Herriott, R.E., Firestone, W.A., 1983. Multisite qualitative policy research: Optimizing description and generalizability. *Educational researcher*, 12(2), 14-19.

Hertogh, C. M., Miesen, B. M., & Eefsting, J. A. (2004). Truth telling and truthfulness in the care for patients with advanced dementia: an ethnographic study in Dutch nursing homes. *Social science & medicine* (1982), 59(8), 1685.

Holmes J., 1999. The detection of psychiatric factors predicting poor outcome in elderly hip fracture patients. MD Thesis. Research School of Medicine, University of Leeds

Holmes, B.J., Best, A., Davies, H., Hunter, D., Kelly, M.P., Marshall, M. and Rycroft-Malone, J., 2017. Mobilising knowledge in complex health systems: a call to action. *Evidence & Policy: A Journal of Research, Debate and Practice*, 13(3), pp.539-560.

House of Commons All-Party Parliamentary Group on Dementia., 2013. *Dementia does not discriminate The experiences of black, Asian and minority ethnic communities* London: The Stationery Office

House of Commons Public Accounts Committee., 2008. *Sixth Report of Session 2007–8: Improving services and support for people with dementia*, London: The Stationery Office.

Inouye S., Bogardus S., Charpentier P., Leo-Summers L., Campora D., Holford T. & Cooney L. (1999) A multicomponent interven- tion to prevent delirium in hospitalized older patients. The New England Journal of Medicine 340, 669–676.

Ishii S, Streim JE, Saliba D (2012) A conceptual framework for rejection of care behaviours: Review of literature and analysis of role of dementia severity. *Journal of the American Directors* Association. 13(1) 11

Jonsson, L., and Wimo, A., 2009. "The cost of dementia in Europe: a review of the evidence, and methodological considerations." *Pharmacoeconomics* 27.5: 391-403.

Kable, A., Guest, M., & McLeod, M. (2012). Resistance to Care: contributing factors and associated behaviours in healthcare facilities. *Journal of advanced nursing*.

Keady J, Jones L. (2010). 'Investigating the causes of behaviours that challenge in people with dementia' *Nursing Older People* 22(9) 25-29

Kitwood T., 1993. Towards a theory of dementia care: the inter- personal process. *Ageing and Society* 13, 51–67.

Knapp, M., Iemmi, V., and Romeo, R., 2012. "Dementia care costs and outcomes: a systematic review." *International journal of geriatric psychiatry*.

Koch, Tamar, and Iliffe. S,. 2010. "Rapid appraisal of barriers to the diagnosis and management of patients with dementia in primary care: a systematic review." *BMC Family Practice* 11.1: 52.

Lakey, L., 2009. *Counting the Costs. Caring for people with dementia on hospital wards.* London: Alzheimer's Society.

Lauder, H., Brown, P. and Halsey, A.H., 2004. Sociology and political arithmetic: some principles of a new policy science1. *The British Journal of Sociology*, 55(1), pp.3-22.

Li H., Melnyk B., McCann R., Chatcheydang J., Koulouglioti C.,

Logsdon, R.G., Teri, L., Weiner, M.F., Gibbons, L.E., Raskind, M., Peskind, E., Koss, E., Thomas, R.G. and Thai, L.J., 1999. Assessment of agitation in Alzheimer's disease: the agitated behavior in dementia scale. *Journal of the American Geriatrics Society*, 47(11), pp.1354-1358.

Lothian, K., Philp, I., 2001. Care of older people: Maintaining the dignity and autonomy of older people in the healthcare setting. *British Medical Journal* 322(7287): 668-670

Maben, J., et al., Exploring the relationship between patients' experiences of care and the influence of staff motivation, affect and wellbeing. *NIHR Service Delivery and Organisation programme*. 2012.

Mays, N., Pope, C. 2000. Qualitative research in health care: Assessing quality in qualitative research. *British Medical Journal*, 320(7226), 50.

Moore, G.F., Audrey, S., Barker, M., Bond, L., Bonell, C., Hardeman, W., Moore, L., O'Cathain, A., Tinati, T., Wight, D. and Baird, J., 2015. Process evaluation of complex interventions: Medical Research Council guidance. *bmj*, 350, p.h1258.

Morrison R.S., Siu A.L., 2000. A Comparison of Pain and Its Treatment in Advanced Dementia and Cognitively Intact Patients with Hip Fracture. *Journal of Pain and Symptom Management* 19(4) 240 - 248

Moyle, W., Olorenshaw, R., Wallis, M., Borbasi, S., 2008. Best practice for the management of older people living with dementia in the acute care setting: a review of the literature. *International Journal of Older People Nursing* 3(2) 121-130.

Mukadam, N. and Sampson, E.L., 2011. A systematic review of the prevalence, associations and outcomes of dementia in older general hospital inpatients. *International Psychogeriatrics*, 23(3), pp.344-355.

National Audit Office, 2007. *Improving services and support for people living with dementia*. London: The Stationery Office.

National Confidential Enquiry into Perioperative Deaths, 2009. Adding Insult to Injury: a Review of the Care of Patients who Died in Hospital with a Primary Diagnosis of Acute Kidney Injury (actute Renal Failure): A Report by the National Confidential Enquiry Into Patient Outcome and Death. National Confidential Enquiry into Patient Outcome and Death.

Newbronner, L., Chamberlain, R., Borthwick, R., Baxter, M., Glendinning, C., 2013. *A Road Less Rocky: Supporting carers of people with dementia* London: Carers Trust

NICE, 2006. The NICE-SCIE guideline on supporting people with dementia and their carers in health and social care. London: The Stationery Office.

Nottingham City and Nottinghamshire County Joint Health Scrutiny Committee 2010

O'Connor CM., Smith R., Nott MT. 2011. 'Using video simulated presence to reduce resistance to care and increase participation of adults with dementia.' *American Journal of Alzheimer's Disease* & Other Dementias 26(4) 317-325.

Patients Association, 2010. *Listen to patients, Speak up for change*. Patients Association.

Patterson, M., et al., 2011. From metrics to meaning: culture change and quality of acute hospital care for older people. NIHR SDO programme project, 3(1501):93

Pawson R, Tilley N. Realistic evaluation. London, UK: Sage; 1997.

Petchey, R., Hughes, J., Pinder, R., Needle, J., Partington, J., & Sims, D., 2013. Allied Health Professionals and management: an ethnographic study. *National Institute for Health Research*.

Phelan, E. A., Borson, S., Grothaus, L., Balch, S., Larson, E. B., 2012. Association of incident dementia with hospitalizations. *Journal of the American Medical Association*, 307(2) 165-172.

Philp, I., 2006. National Service Framework for Older People London: DoH

Pinkert, C., Holle, B., 2012. People with dementia in acute hospitals: Literature review of prevalence and reasons for hospital admission, Zeitschrift fur Gerontologie und Geriatrie 45(8):728-34

Pitfield, C., Shahriyarmolki, K., and Livingston. G., 2011. "A systematic review of stress in staff caring for people with dementia living in 24-hour care settings." *International Psychogeriatrics* 23.1: 4.

Pizzacalla A., Montemuro M., Coker E., Schindel M.L., Gillies L., Robinson K., Pepper H., Benner J., Gusciora J. 2015. Gentle Persuasive Approaches: Introducing an Educational Program on an Orthopaedic Unit for Staff Caring for Patients With Dementia and Delirium. *Orthopaedic Nursing* 34(2) 101-107

Popay, J., & Williams, G. 1998. Qualitative research and evidence-based healthcare. *Journal of the Royal Society of Medicine*, 91(Suppl 35), 32.

Quinlan, E., 2009. The 'actualities' of knowledge work: an institutional ethnography of multidisciplinary primary health careteams. *Sociology of health & illness*, 31(5):625-641

Radar, J., Doan, J. and Schwab, M., 1985. How to decrease wandering, a form of agenda behaviour. *Geriatric Nursing*, 6(4), pp.196-199.

Rantala M., Kankkunen P., Kvist T., Hartikainen S. 2014. Barriers to postoperative pain management in hip fracture patients with dementia as evaluated by nursing staff. *Pain Management Nursing* 15(1) 208-19

Rockwood, K., & Hubbard, R. 2004. Frailty and the geriatrician. Age and Ageing, 33(5), 429-430.

Rosaldo, R. (1989). Culture and truth: the remaking of social analysis. Boston: Beacon.

Royal College of Psychiatrists, 2006. Who Cares Wins: Improving the Outcome for Older People Admitted to the General Hospital London: RCP

Royal College of Psychiatrists, 2013. *National Audit of Dementia care in general hospitals 2012-13:* Second round audit report and update. London: HQIP.

Russ, C. T., Shenkin S.D., Reynish. E., Ryan, T., Anderson, D., MacLullich, A.M.J., 2012. Dementia in acute hospital inpatients: the role of the geriatrician, *Age and Ageing*, Volume 41(3), 282–284

Sager M.A., Franke T., Inouye SK. 1996. Functional Outcomes of Acute Medical Illness and Hospitalization in Older Persons. *Arch Intern Med.* 156(6):645-652.

Sampson, E.L., 2006 "Differences in care received by patients with and without dementia who died during acute hospital admission: A retrospective case note study." *Age and Ageing* 35.2: 187-189.

Sampson, E. L., Blanchard, M. R., Jones, L., Tookman, A., & King, M., 2009. Dementia in the acute hospital: prospective cohort study of prevalence and mortality. *The British Journal of Psychiatry*, 195(1):61-66.

Sampson, E.L., Leurent, B., Blanchard, M.R., Jones, L. and King, M., 2013. Survival of people with dementia after unplanned acute hospital admission: a prospective cohort study. *International journal of geriatric psychiatry*, 28(10), pp.1015-1022.

Sampson, E.L., White, N., Lord, K., Leurent, B., Vickerstaff, V., Scott, S. and Jones, L., 2015. Pain, agitation, and behavioural problems in people with dementia admitted to general hospital wards: a longitudinal cohort study. *Pain*, 156(4), p.675.

Schoenmakers, B., Buntinx, F., and Jan Delepeleire, J., 2010. "Factors determining the impact of care-giving on caregivers of elderly patients with dementia. A systematic literature review." *Maturitas* 66.2: 191-200.

Scottish Government, 2010. *Scotland's National Dementia Strategy*. Edinburgh : Scottish Government.

Seitz, D., Prorok, J., & Horgan, S. (2012). Health care experiences of persons with dementia and their caregivers: A meta-ethnography of qualitative studies. *Alzheimer's & Dementia*, 8(4), P562.

Starr, S.L. 1999. The ethnography of infrastructure. American behavioral scientist, 43(3), 377-391.

Stevenson, G.S., Khan, M.A. and Perumal, N., 2009. A psychiatric intensive care unit for older adults: an interval comparison of admissions. *International psychogeriatrics*, 21(2), pp.278-285.

Suddaby, R. (2006). From the editors: What grounded theory is not. *Academy of management journal*, 49(4), 633-642.

Tadd, W., Hillman, A., Calnan, M., Calnan, S., Read, S., & Bayer, A. (2012). From Right place—Wrong person, to Right place—Right person: dignified care for older people. *Journal of Health Services Research & Policy*, 17(suppl 2), 30-36.

Tadd, W., Hillman, A., Calnan, S., Calnan, M., Bayer, T., & Read, S. (2011). 'Right place-wrong person: dignity in the acute care of older people.' *Quality in Ageing and Older Adults*, 12(1), 33-43.

The House of Lords House of Commons Joint Committee on Human Rights., 2006 *The Human Rights of Older People in Healthcare Eighteenth Report of Session 2006-07 Volume I - Report and Formal Minutes* HL 156-I, HC 378-I 2006.

Thornlow, D.K., Anderson, R., and Oddone, E. 2009. Cascade iatrogenesis: Factors leading to the development of adverse events in hospitalized older adults. *International Journal of Nursing Studies* 46(11) 1528–1535

Tolson, D., Smith, M., Knight, P., 1999. An investigation of the components of best nursing practice in the care of acutely ill hospitalized older patients with coincidental dementia: a multimethod design. *Journal of Advanced Nursing* 30(5) 1127-1136.

Torti Jr, F. M., Gwyther, L. P., Reed, S. D., Friedman, J. Y., & Schulman, K. A. (2004). A multinational review of recent trends and reports in dementia caregiver burden. *Alzheimer Disease & Associated Disorders*, 18(2), 99-109.

Traynor, V., Inoue. K., Crookes. P., 2011. "Literature review: understanding nursing competence in dementia care." *Journal of Clinical Nursing* 20.13-14: 1948-1960.

van der Geest S, Finkler K. 2004 Hospital ethnography: introduction. Soc Sci Med. 59:1995-2001.

Van Maanen, J. (2011). Tales of the field: On writing ethnography. University of Chicago Press.

Vasse, E., et al., 2010. "A systematic review of communication strategies for people with dementia in residential and nursing homes." *International Psychogeriatrics* 22.2: 189.

Vaughan, D., 2005. "On the relevance of ethnography for the production of public sociology and policy." *The British journal of sociology* 56.3: 411-416.

Vizard, P. and Burchardt, T., 2015. Older people's experiences of dignity and nutrition during hospital stays: secondary data analysis using the Adult Inpatient Survey.

Vogt, F. 2002. No ethnography without comparison: the methodological significance of comparison in ethnographic research. *Studies in Educational Ethnography*, 6, 23-42.

Wasson K et al., 2001. 'Food refusal and dysphagia in older people with dementia: ethical and practical issues' *Palliative Nursing* 7 (10) 463-471

Watkin, L., Blanchard, M.R., Tookman, A. and Sampson, E.L., 2012. Prospective cohort study of adverse events in older people admitted to the acute general hospital: risk factors and the impact of dementia. *International journal of geriatric psychiatry*, 27(1), pp.76-82.

Werner P., Tabak N., Alpert R., Bergman R. 2002. Interventions used by nursing staff members with psychogeriatric patients resisting care. *International Journal of Nursing Studies* 39 461-467

Williamson, S., Twelvetree, T., Thompson, J., & Beaver, K. (2012). An ethnographic study exploring the role of ward-based Advanced Nurse Practitioners in an acute medical setting. Journal of advanced nursing, 68(7), 1579-1588.

Ybema, S., Yanow, D., Wels, H., & Kamsteeg, F. H. (Eds.). 2009. *Organizational ethnography:* Studying the complexity of everyday life. SAGE Publications Limited.